

Chiral and Deconfinement Aspects of (2+1)-flavor QCD

Bernd-Jochen Schaefer

Karl-Franzens-Universität Graz, Austria



17th - 23rd January, 2010

XXXVIII Hirschepp Workshop

Strongly Interacting Matter under Extreme Conditions

Hirschepp, Austria

QCD Phase Transitions

QCD: two phase transitions:


- restoration of chiral symmetry

$$SU_{L+R}(N_f) \rightarrow SU_L(N_f) \times SU_R(N_f)$$

order parameter:

$$\langle \bar{q}q \rangle \begin{cases} > 0 \Leftrightarrow \text{symmetry broken, } T < T_c \\ = 0 \Leftrightarrow \text{symmetric phase, } T > T_c \end{cases}$$

associate limit: $m_q \rightarrow 0$



chiral transition: spontaneous restoration of global $SU_L(N_f) \times SU_R(N_f)$ at high T

QCD Phase Transitions

QCD: two phase transitions:

- 1 restoration of chiral symmetry
- 2 de/confinement (center symmetry)


order parameter:

$$\Phi \begin{cases} = 0 \Leftrightarrow \text{confined phase}, & T < T_c \\ > 0 \Leftrightarrow \text{deconfined phase}, & T > T_c \end{cases}$$

$$\Phi = \frac{1}{N_c} \langle \text{tr}_c \mathcal{P} e^{i \int_0^\beta d\tau A_0(\tau, \vec{x})} \rangle$$

associate limit: $m_q \rightarrow \infty$

→ related to free energy of a static quark state: $\Phi = e^{-\beta F_q}$




QCD Phase Transitions

QCD: two phase transitions:

- 1 restoration of chiral symmetry
- 2 de/confinement (center symmetry)

order parameter:


$$\Phi \left\{ \begin{array}{l} = 0 \Leftrightarrow \text{confined phase}, \quad T < T_c \\ > 0 \Leftrightarrow \text{deconfined phase}, \quad T > T_c \end{array} \right.$$



alternative:

- dressed Polyakov loop (or dual condensate)
it relates chiral and deconfinement transition to spectral properties of Dirac operator

The conjectured QCD Phase Diagram



At densities/temperatures of interest
only model calculations available

Open issues:

related to chiral & deconfinement transition

- ▷ existence of CEP?
- ▷ its location?
- ▷ additional CEPs?
How many?
- ▷ coincidence of both transitions at $\mu = 0$?
- ▷ quarkyonic phase at $\mu > 0$?
- ▷ chiral CEP/
deconfinement CEP?
- ▷ so far only MFA results
effect of fluctuations (e.g. size of crit. reg.)?
- ▷ ...

effective models:

- 1 Quark-meson model (renormalizable)
- 2 Polyakov–quark-meson model

or other models e.g. NJL

or PNJL models


Charts of QCD Critical End Points

model studies vs. lattice simulations

Black points: models


Lines & green points: lattice

Red points: Freezeout points for HIC



Phase diagram in (T, μ_B, m_q) -space


Chiral limit: $(m_q = 0)$ $SU(2) \times SU(2) \sim O(4)$ -symmetry \longrightarrow 4 modes critical $\sigma, \vec{\pi}$



Phase diagram in (T, μ_B, m_q) -space

Chiral limit: $(m_q = 0)$ $SU(2) \times SU(2) \sim O(4)$ -symmetry \longrightarrow 4 modes critical $\sigma, \vec{\pi}$

$m_q \neq 0$: no symmetry remains \longrightarrow only one critical mode σ (**Ising**) ($\vec{\pi}$ massive)



General properties

- chiral limit
tricritical point
(Gaussian fixed point)
- finite m_q
critical endpoints
(3D-Ising class)

Outline

- **Three-Flavor Quark-Meson Model**
- ...with Polyakov loop dynamics
- Finite density extrapolations

$N_f = 3$ Quark-Meson (QM) model

- Model Lagrangian: $\mathcal{L}_{\text{qm}} = \mathcal{L}_{\text{quark}} + \mathcal{L}_{\text{meson}}$

Quark part with Yukawa coupling g:

$$\mathcal{L}_{\text{quark}} = \bar{q}(i\partial - g \frac{\lambda_a}{2}(\sigma_a + i\gamma_5 \pi_a))q$$

Meson part: scalar σ_a and pseudoscalar π_a nonet

fields: $\phi = \sum_{a=0}^8 \frac{\lambda_a}{2}(\sigma_a + i\pi_a)$

$$\begin{aligned}\mathcal{L}_{\text{meson}} = & \text{tr}[\partial_\mu \phi^\dagger \partial^\mu \phi] - m^2 \text{tr}[\phi^\dagger \phi] - \lambda_1 (\text{tr}[\phi^\dagger \phi])^2 - \lambda_2 \text{tr}[(\phi^\dagger \phi)^2] + c [\det(\phi) + \det(\phi^\dagger)] \\ & + \text{tr}[H(\phi + \phi^\dagger)]\end{aligned}$$

- explicit symmetry breaking matrix: $H = \sum_a \frac{\lambda_a}{2} h_a$
- $U(1)_A$ symmetry breaking implemented by 't Hooft interaction

Phase diagram for $N_f = 2 + 1$ ($\mu \equiv \mu_q = \mu_s$)


- Model parameter fitted to (pseudo)scalar meson spectrum:
- PDG: $f_0(600)$ mass=(400 . . . 1200) MeV → broad resonance

→ existence of CEP depends on m_σ !

Example: $m_\sigma = 600$ MeV (lower lines), 800 and 900 MeV (here mean-field approximation)

with $U(1)_A$



[BJS, M. Wagner '09]



In-medium meson masses

Finite temperature axis: $\mu = 0$

masses without $U(1)_A$ anomaly





- At low temperatures: mesons dominate
- At high temperatures: quarks dominate

In-medium meson masses

Finite temperature axis: $\mu = 0$

masses with $U(1)_A$ anomaly





- At low temperatures: mesons dominate
- At high temperatures: quarks dominate

In-medium meson masses

slide through CEP: $\mu = \mu_c$

masses with $U(1)_A$ anomaly



- At low temperatures: mesons dominate
- At high temperatures: quarks dominate


Mass sensitivity

Chiral limit: RG arguments → for $N_f \geq 3$ first-order

[Pisarski, Wilczek '84]

Columbia plot:

[Brown et al. '90]



Mass sensitivity

Chiral limit: RG arguments → for $N_f \geq 3$ first-order



[Pisarski, Wilczek '84]

- variation of m_π and m_K :

$m_\pi/m_\pi^* = 0.49$ (lower line), $0.6, 0.8 \dots, 1.36$ (upper line)


$m_\pi^* = 138 \text{ MeV}$, $m_K^* = 496 \text{ MeV}$, fixed ratio $m_\pi/m_K = m_\pi^*/m_K^*$

with $U(1)_A$, $m_\sigma = 800 \text{ MeV}$




Mass sensitivity (lattice, $N_f = 3$, $\mu_B \neq 0$)

Standard scenario: $m_c(\mu)$ increasing



Nonstandard scenario: $m_c(\mu)$ decreasing




[de Forcrand, Philipsen: hep-lat/0611027]


Chiral critical surface ($m_\sigma = 800$ MeV)

→ standard scenario for $m_\sigma = 800$ MeV (as expected)

with $U(1)_A$



without $U(1)_A$



[BJS, M. Wagner, '09]


Note: 't Hooft coupling μ -independent
PNJL with (unrealistic) large vector int. → bending of surface

Chiral critical surface for different m_σ


▷ CEP vanishes for $m_\sigma > 800$ MeV → **non-standard scenario possible?**

No → three cuts of critical surface along fixed m_π/m_K ratio through physical point

critical μ_c



critical T_c



[BJS, M. Wagner, '09]

Outline

- Three-Flavor Quark-Meson Model
- ...with Polyakov loop dynamics
- Finite density extrapolations

Polyakov-quark-meson (PQM) model

- Lagrangian $\mathcal{L}_{\text{PQM}} = \mathcal{L}_{\text{qm}} + \mathcal{L}_{\text{pol}}$ with $\mathcal{L}_{\text{pol}} = -\bar{q}\gamma_0 A_0 q - \mathcal{U}(\phi, \bar{\phi})$

- polynomial Polyakov loop potential:

Polyakov 1978, Meisinger 1996, Pisarski 2000

$$\frac{\mathcal{U}(\phi, \bar{\phi})}{T^4} = -\frac{b_2(\textcolor{red}{T}, \textcolor{red}{T}_0)}{2} \phi \bar{\phi} - \frac{b_3}{6} (\phi^3 + \bar{\phi}^3) + \frac{b_4}{16} (\phi \bar{\phi})^2$$

with

$$b_2(T, T_0) = a_0 + a_1(T_0/T) + a_2(T_0/T)^2 + a_3(T_0/T)^3$$

Polyakov-quark-meson (PQM) model

■ Lagrangian $\mathcal{L}_{\text{PQM}} = \mathcal{L}_{\text{qm}} + \mathcal{L}_{\text{pol}}$ with $\mathcal{L}_{\text{pol}} = -\bar{q}\gamma_0 A_0 q - \mathcal{U}(\phi, \bar{\phi})$

■ polynomial Polyakov loop potential:

Polyakov 1978, Meisinger 1996, Pisarski 2000

$$\frac{\mathcal{U}(\phi, \bar{\phi})}{T^4} = -\frac{b_2(T, T_0)}{2} \phi \bar{\phi} - \frac{b_3}{6} (\phi^3 + \bar{\phi}^3) + \frac{b_4}{16} (\phi \bar{\phi})^2$$

with

$$b_2(T, T_0) = a_0 + a_1(T_0/T) + a_2(T_0/T)^2 + a_3(T_0/T)^3$$

■ logarithmic potential:

Rößner et al. 2007

$$\frac{\mathcal{U}_{\log}}{T^4} = -\frac{1}{2} a(T) \bar{\phi} \phi + b(T) \ln \left[1 - 6\bar{\phi} \phi + 4 (\phi^3 + \bar{\phi}^3) - 3 (\bar{\phi} \phi)^2 \right]$$

with

$$a(T) = a_0 + a_1(T_0/T) + a_2(T_0/T)^2 \quad \text{and} \quad b(T) = b_3(T_0/T)^3$$

Polyakov-quark-meson (PQM) model

■ Lagrangian $\mathcal{L}_{\text{PQM}} = \mathcal{L}_{\text{qm}} + \mathcal{L}_{\text{pol}}$ with $\mathcal{L}_{\text{pol}} = -\bar{q}\gamma_0 A_0 q - \mathcal{U}(\phi, \bar{\phi})$

■ polynomial Polyakov loop potential:

Polyakov 1978, Meisinger 1996, Pisarski 2000

$$\frac{\mathcal{U}(\phi, \bar{\phi})}{T^4} = -\frac{b_2(T, T_0)}{2} \phi \bar{\phi} - \frac{b_3}{6} (\phi^3 + \bar{\phi}^3) + \frac{b_4}{16} (\phi \bar{\phi})^2$$

with

$$b_2(T, T_0) = a_0 + a_1(T_0/T) + a_2(T_0/T)^2 + a_3(T_0/T)^3$$

■ Fukushima

Fukushima 2008

$$\mathcal{U}_{\text{Fuku}} = -bT \left\{ 54e^{-a/T} \phi \bar{\phi} + \ln \left[1 - 6\bar{\phi}\phi + 4(\phi^3 + \bar{\phi}^3) - 3(\bar{\phi}\phi)^2 \right] \right\}$$

with

a controls deconfinement b strength of mixing chiral & deconfinement

Polyakov-quark-meson (PQM) model

■ Lagrangian $\mathcal{L}_{\text{PQM}} = \mathcal{L}_{\text{qm}} + \mathcal{L}_{\text{pol}}$ with $\mathcal{L}_{\text{pol}} = -\bar{q}\gamma_0 A_0 q - \mathcal{U}(\phi, \bar{\phi})$

■ polynomial Polyakov loop potential:

Polyakov 1978, Meisinger 1996, Pisarski 2000

$$\frac{\mathcal{U}(\phi, \bar{\phi})}{T^4} = -\frac{b_2(\textcolor{red}{T}, \textcolor{red}{T}_0)}{2} \phi \bar{\phi} - \frac{b_3}{6} (\phi^3 + \bar{\phi}^3) + \frac{b_4}{16} (\phi \bar{\phi})^2$$

with

$$b_2(T, T_0) = a_0 + a_1(T_0/T) + a_2(T_0/T)^2 + a_3(T_0/T)^3$$

in presence of dynamical quarks: $T_0 = T_0(\textcolor{red}{N_f}, \mu)$

BJS, Pawłowski, Wambach, 2007

N_f		0	1	2	2 + 1	3
T_0 [MeV]		270	240	208	187	178

$\mu \neq 0$: $\bar{\phi} > \phi$


since $\bar{\phi}$ is related to free energy gain of antiquarks

in medium with more quarks \rightarrow antiquarks are more easily screened.


Finite temperature and finite μ (PQM $N_f = 2$)

without $T_0(\mu)$ -modifications in Polyakov loop potential:

order parameters



[BJS, Pawłowski, Wambach '07]




Phase diagrams $N_f = 2$

[BJS, Pawłowski, Wambach '07]

in mean field approximation
chiral transition and 'deconfinement' coincide


■ for PQM model
 $N_f = 2$



Phase diagrams $N_f = 2$

[BJS, Pawłowski, Wambach '07]


in mean field approximation
chiral transition and 'deconfinement' coincide



Phase diagrams $N_f = 2$

[BJS, Pawłowski, Wambach '07]

in mean field approximation
chiral transition and 'deconfinement' coincide



- for PQM model
 $N_f = 2$
- for PQM model
 $N_f = 2$
with
 $T_0(\mu)$ -modification
in Polyakov loop
potential
(lower lines)

Phase diagram $N_f = 2 + 1$


[BJS, M. Wagner; in preparation '10]

influence of Polyakov loop


Logarithmic Polyakov loop potential

Mean-field approximation

$T_0 = 270 \text{ MeV}$ (constant)



$T_0(\mu)$ (i.e. with μ corrections)





shrinking of possible quarkyonic phase

Critical region

contour plot of **size of the critical region** around CEP

defined via fixed ratio of susceptibilities: $R = \chi_q / \chi_q^{\text{free}}$

→ compressed with Polyakov loop



[BJS, M. Wagner; in preparation '10]


Critical region

similar conclusion if **fluctuations** are included

fluctuations via Renormalization Group

comparison: $N_f = 2$ QM model

Mean Field \leftrightarrow RG analysis



[BJS, J. Wambach '06]



Isentropes $s/n = \text{const}$ and Focussing

[E. Nakano, BJS, B.Stokic, B.Friman, K.Redlich '10]

here: $N_f = 2$ QM model: kink in MFA are washed out in FRG

→ no focussing if fluctuations taken into account

a) influence of Dirac term b) smallness of critical region





Isentropes $s/n = \text{const}$ and Focussing

[E. Nakano, BJS, B.Stokic, B.Friman, K.Redlich '10]

here: $N_f = 2$ QM model: kink in MFA are washed out in FRG

→ no focussing if fluctuations taken into account

a) influence of Dirac term b) smallness of critical region



Isentropes $s/n = \text{const}$ and Focussing

[E. Nakano, BJS, B.Stokic, B.Friman, K.Redlich '10]

here: $N_f = 2$ QM model: kink in MFA are washed out in FRG

→ no focussing if fluctuations taken into account

- a) influence of Dirac term
- b) smallness of crit region

kink structure at boundary in mean field approximation

⇒ remnant of first-order transition in chiral limit


if Dirac term neglected

QCD Thermodynamics $N_f = 2 + 1$

[BJS, M. Wagner, J. Wambach; arXiv:0910.5628]

$$\text{SB limit: } \frac{p_{\text{SB}}}{T^4} = 2(N_c^2 - 1) \frac{\pi^2}{90} + N_f N_c \frac{7\pi^2}{180}$$

(P)QM models (three different Polyakov loop potentials) versus QCD lattice simulations



- ▷ solid lines:
PQM with lattice masses
(HotQCD)
- ▷ dashed lines:
(P)QM with realistic masses



lattice data: [Bazavov et al. '09]

QCD Thermodynamics $N_f = 2 + 1$

[BJS, M. Wagner, J. Wambach; arXiv:0910.5628]

$$\text{SB limit: } \frac{P_{\text{SB}}}{T^4} = 2(N_c^2 - 1) \frac{\pi^2}{90} + N_f N_c \frac{7\pi^2}{180}$$

(P)QM models (three different Polyakov loop potentials) versus QCD lattice simulations



solid lines: $m_\pi \sim 220, m_K \sim 503$ MeV (HotQCD)

[Bazavov et al. '09]

Outline

- Three-Flavor Quark-Meson Model
- ...with Polyakov loop dynamics
- **Finite density extrapolations**

Finite density extrapolations $N_f = 2 + 1$

Taylor expansion:

$$\frac{p(T, \mu)}{T^4} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu}{T}\right)^n \quad \text{with} \quad c_n(T) = \frac{1}{n!} \frac{\partial^n (p(T, \mu)/T^4)}{\partial (\mu/T)^n} \Big|_{\mu=0}$$

high temperature limits:

$$c_0(T \rightarrow \infty) = \frac{7N_c N_f \pi^2}{180},$$

$$c_2(T \rightarrow \infty) = \frac{N_c N_f}{6},$$


$$c_4(T \rightarrow \infty) = \frac{N_c N_f}{12\pi^2}$$

$$c_n(T \rightarrow \infty) = 0 \text{ for } n > 4.$$

Finite density extrapolations $N_f = 2 + 1$

Taylor expansion:

$$\frac{p(T, \mu)}{T^4} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu}{T}\right)^n \quad \text{with} \quad c_n(T) = \frac{1}{n!} \frac{\partial^n (p(T, \mu)/T^4)}{\partial (\mu/T)^n} \Big|_{\mu=0}$$



convergence radii:

limited by first-order line?

$$\rho_{2n} = \left| \frac{c_2}{c_{2n}} \right|^{1/(2n-2)}$$

[C. Schmidt '09]



Finite density extrapolations $N_f = 2 + 1$

Taylor expansion:

$$\frac{p(T, \mu)}{T^4} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu}{T}\right)^n \quad \text{with} \quad c_n(T) = \frac{1}{n!} \frac{\partial^n (p(T, \mu)/T^4)}{\partial (\mu/T)^n} \Big|_{\mu=0}$$

first three coefficients:


c_0 : pressure at $\mu = 0$



Finite density extrapolations $N_f = 2 + 1$


Taylor expansion:

$$\frac{p(T, \mu)}{T^4} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu}{T} \right)^n \quad \text{with} \quad c_n(T) = \frac{1}{n!} \frac{\partial^n (p(T, \mu)/T^4)}{\partial (\mu/T)^n} \Bigg|_{\mu=0}$$




[Miao et al. '08]

Taylor coefficients c_n numerically known to high order, e.g. $n = 22$



Finite density extrapolations $N_f = 2 + 1$




- ▷ this technique applied to PQM model
- ▷ investigation of convergence properties of Taylor series
- ▷ properties of c_n
 - oscillating
 - increasing amplitude
 - no numerical noise
 - small outside transition region
 - number of roots increasing
 - 26th order

[F. Karsch, BJS, M. Wagner, J. Wambach; in preparation '10]


Can we locate the QCD critical endpoint with the Taylor expansion ?

Susceptibility $N_f = 2 + 1$ PQM model


$$\mu/T = 0.8$$



$$\mu/T = \mu_c/T_c$$




$$\mu/T = 3.$$




Susceptibility $N_f = 2 + 1$ PQM model


$\mu/T = 0.8$




$\mu/T = \mu_c/T_c$



$\mu/T = 3.$



convergence radius



Susceptibility $N_f = 2 + 1$ PQM model

Findings:


- simply Taylor expansion: slow convergence
high orders needed
disadvantage for lattice simulations
- Taylor applicable within convergence radius
also for $\mu/T > 1$
- but 1st order transition not resolvable
expansion around $\mu = 0$

Summary

- $N_f = 2$ and $N_f = 2 + 1$ chiral (Polyakov)-quark-meson model study
 - Mean-field approximation and FRG
 - with and without axial anomaly
- novel AD technique: high order Taylor coefficients, here: $n = 26$

Findings:

- ▷ Parameter in Polyakov loop potential:
 $T_0 \Rightarrow T_0(N_f, \mu)$
- ▷ Chiral & deconfinement transition possibly **coincide** for $N_f = 2$ with $T_0(\mu)$ -corrections but possibly not for $N_f = 2 + 1$
- ▷ Mean-field approximation encouraging but effects of Dirac term point to interesting physics if fluctuations are considered
→ FRG with PQM truncation
- ▷ Taylor coefficient $c_n(T) \rightarrow$ **high order**
- ⇒ **convergence properties** of Taylor expansion



Outlook:

- include glue dynamics with FRG → full QCD