

Hirscheegg 2013: Astrophysics and Nuclear Structure
Hirscheegg, Austria, January 26–February 1, 2013

Explosion Models of Core- Collapse Supernovae Status of Modeling at Garching

Hans-Thomas Janka
Max Planck Institute for Astrophysics, Garching

Supernova and neutron star merger research is team effort

Students & postdocs:



Alexandra Gessner
Robert Bollig
Andreas Voth

Thomas Ertl
Tobias Melson
Florian Hanke
Lorenz Hüdepohl
Janina von Groote
Else Pllumbi

Andreas Bauswein
Oliver Just
Pedro Montero
Bernhard Müller
Marcella Ugliano
Annop Wongwathanarat



Heinzi-Ado Arnolds (MPA, 2012)



Collaborators:

Ewald Müller
Martin Obergaulinger
Andreas Marek
Stephane Goriely
Nick Stergioulas
Thomas Baumgarte
Georg Raffelt
Victor Utrobin
Shinya Wanajo

Outline

- **Introduction to core-collapse supernova dynamics**
- **The neutrino-driven mechanism**
- **Status of self-consistent models in two dimensions**
- **The dimension conundrum: How does 3D differ from 2D?**

Explosion Mechanism
by
Neutrino Heating


Explosion:

Shock wave expands into outer stellar layers, heats and ejects them.


Creation of radioactive nickel in shock-heated Si-layer.

Shock wave

Proto-neutron star (PNS)



Neutrino Heating and Cooling



- Neutrino heating:

$$q_\nu^+ = 1.544 \times 10^{20} \left(\frac{L_{\nu_e}}{10^{52} \text{ erg s}^{-1}} \right) \left(\frac{T_{\nu_e}}{4 \text{ MeV}} \right)^2 \times \left(\frac{100 \text{ km}}{r} \right)^2 (Y_n + Y_p) \quad \left[\frac{\text{erg}}{\text{g s}} \right]$$

- Neutrino cooling:

$$C = 1.399 \times 10^{20} \left(\frac{T}{2 \text{ MeV}} \right)^6 (Y_n + Y_p) \quad \left[\frac{\text{erg}}{\text{g s}} \right]$$

$$Q_\nu^+ = q_\nu^+ M_g$$

$$\sim 9.4 \times 10^{51} \frac{\text{erg}}{\text{s}} \left(\frac{k_B T_\nu}{4 \text{ MeV}} \right)^2 \left(\frac{L_\nu}{3 \cdot 10^{52} \text{ erg/s}} \right) \left(\frac{M_g}{0.01 M_\odot} \right) \left(\frac{R_g}{100 \text{ km}} \right)^{-2}$$

$$E_N \sim Q_\nu^+ t_{\text{dwell}}$$

$$\sim 9.4 \times 10^{50} \text{ erg} \left(\frac{k_B T_\nu}{4 \text{ MeV}} \right)^2 \left(\frac{L_\nu}{3 \cdot 10^{52} \text{ erg/s}} \right) \times \left(\frac{M_g}{0.01 M_\odot} \right)^2 \left(\frac{\dot{M}}{0.1 M_\odot \text{ s}^{-1}} \right)^{-1} \left(\frac{R_g}{100 \text{ km}} \right)^{-2}$$

$$t_{\text{dwell}} \approx \frac{M_g}{\dot{M}}$$


Hydrodynamic instabilities

1D-2D Differences in Parametric Explosion Models

- Nordhaus et al. (ApJ 720 (2010) 694) and Murphy & Burrows (2008) performed 1D & 2D simulations with **simple neutrino- heating and cooling terms (no neutrino transport but lightbulb)** and found up to ~30% improvement in 2D for 15 M_{sun} progenitor star.

$$\mathcal{H} = 1.544 \times 10^{20} \left(\frac{L_{\nu_e}}{10^{52} \text{ erg s}^{-1}} \right) \left(\frac{T_{\nu_e}}{4 \text{ MeV}} \right)^2 \times \left(\frac{100 \text{ km}}{r} \right)^2 (Y_n + Y_p) e^{-\tau_{\nu_e}} \left[\frac{\text{erg}}{\text{g s}} \right]$$

$$\mathcal{C} = 1.399 \times 10^{20} \left(\frac{T}{2 \text{ MeV}} \right)^6 (Y_n + Y_p) e^{-\tau_{\nu_e}} \left[\frac{\text{erg}}{\text{g s}} \right]$$



But: Is neutrino heating strong enough to initiate the explosion?

Most sophisticated, self-consistent numerical simulations of the explosion mechanism in 2D and 3D are necessary!

Predictions of Signals from SN Core

hydrodynamics of stellar plasma

Relativistic gravity

(nuclear) EoS

neutrino physics

progenitor conditions

SN explosion models

neutrinos

LC, spectra

nucleosynthesis

gravitational waves

explosion asymmetries,
pulsar kicks

explosion energies, remnant masses

Explosion Mechanism:
Most Sophisticated Current
Models

General-Relativistic 2D Supernova Models of the Garching Group

(Müller B., PhD Thesis (2009);
Müller et al., ApJS, (2010))

GR hydrodynamics (CoCoNuT)

$$\frac{\partial\sqrt{\gamma\rho}W}{\partial t} + \frac{\partial\sqrt{-g\rho}W\hat{v}^i}{\partial x^i} = 0, \quad (2.5)$$

$$\frac{\partial\sqrt{\gamma\rho h}W^2v_j}{\partial t} + \frac{\partial\sqrt{-g}\left(\rho hW^2v_j\hat{v}^i + \delta_j^i P\right)}{\partial x^i} = \frac{1}{2}\sqrt{-g}T^{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial x^j} + \left(\frac{\partial\sqrt{\gamma}S_j}{\partial t}\right)_C, \quad (2.6)$$

$$\frac{\partial\sqrt{\gamma}\tau}{\partial t} + \frac{\partial\sqrt{-g}\left(\tau\hat{v}^i + Pv^i\right)}{\partial x^i} = \alpha\sqrt{-g}\left(T^{\mu 0}\frac{\partial\ln\alpha}{\partial x^\mu} - T^{\mu\nu}\Gamma_{\mu\nu}^0\right) + \left(\frac{\partial\sqrt{\gamma}\tau}{\partial t}\right)_C. \quad (2.7)$$

$$\frac{\partial\sqrt{\gamma\rho}WY_e}{\partial t} + \frac{\partial\sqrt{-g\rho}WY_e\hat{v}^i}{\partial x^i} = \left(\frac{\partial\sqrt{\gamma\rho}WY_e}{\partial t}\right)_C, \quad (2.8)$$

$$\frac{\partial\sqrt{\gamma\rho}WX_k}{\partial t} + \frac{\partial\sqrt{-g\rho}WX_k\hat{v}^i}{\partial x^i} = 0. \quad (2.9)$$

CFC metric equations

$$\hat{\Delta}\Phi = -2\pi\phi^5\left(E + \frac{K_{ij}K^{ij}}{16\pi}\right), \quad (2.10)$$

$$\hat{\Delta}(\alpha\Phi) = 2\pi\alpha\phi^5\left(E + 2S + \frac{7K_{ij}K^{ij}}{16\pi}\right), \quad (2.11)$$

$$\hat{\Delta}\beta^i = 16\pi\alpha\phi^4S^i + 2\phi^{10}K^{ij}\hat{\nabla}_j\left(\frac{\alpha}{\Phi^6}\right) - \frac{1}{3}\hat{\nabla}^i\hat{\nabla}_j\beta^j, \quad (2.12)$$

$$\begin{aligned} & \frac{\partial W(\hat{J} + v_r\hat{H})}{\partial t} + \frac{\partial}{\partial r}\left[\left(W\frac{\alpha}{\phi^2} - \beta_r v_r\right)\hat{H} + \left(Wv_r\frac{\alpha}{\phi^2} - \beta_r\right)\hat{J}\right] - \\ & \frac{\partial}{\partial \varepsilon}\left\{W\varepsilon\hat{J}\left[\frac{1}{r}\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right) + 2\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right)\frac{\partial\ln\phi}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right] + \right. \\ & W\varepsilon\hat{H}\left[v_r\left(\frac{\partial\beta_r\phi^2}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right) - \frac{\alpha}{\phi^2}\frac{\partial\ln\alpha W}{\partial r} + \alpha W^2\left(\beta_r\frac{\partial v_r}{\partial r} - \frac{\partial v_r}{\partial t}\right)\right] - \\ & \left.\varepsilon\hat{K}\left[\frac{\beta_r W}{r} - \frac{\partial\beta_r W}{\partial r} + Wv_{r,r}\frac{\partial}{\partial r}\left(\frac{\alpha}{r\phi^2}\right) + W^3\left(\frac{\alpha}{\phi^2}\frac{\partial v_r}{\partial r} + v_r\frac{\partial v_r}{\partial t}\right)\right]\right\} - \\ & W\hat{J}\left[\frac{1}{r}\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right) + 2\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right)\frac{\partial\ln\phi}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right] - \\ & W\hat{H}\left[v_r\left(\frac{\partial\beta_r\phi^2}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right) - \frac{\alpha}{\phi^2}\frac{\partial\ln\alpha W}{\partial r} + \alpha W^2\left(\beta_r\frac{\partial v_r}{\partial r} - \frac{\partial v_r}{\partial t}\right)\right] + \\ & \hat{K}\left[\frac{\beta_r W}{r} - \frac{\partial\beta_r W}{\partial r} + Wv_{r,r}\frac{\partial}{\partial r}\left(\frac{\alpha}{r\phi^2}\right) + W^3\left(\frac{\alpha}{\phi^2}\frac{\partial v_r}{\partial r} + v_r\frac{\partial v_r}{\partial t}\right)\right] = \alpha\hat{C}^{(0)}, \end{aligned} \quad (2.28)$$

Neutrino transport (VERTEX)

$$\begin{aligned} & \frac{\partial W(\hat{H} + v_r\hat{K})}{\partial t} + \frac{\partial}{\partial r}\left[\left(W\frac{\alpha}{\phi^2} - \beta_r v_r\right)\hat{K} + \left(Wv_r\frac{\alpha}{\phi^2} - \beta_r\right)\hat{H}\right] - \\ & \frac{\partial}{\partial \varepsilon}\left\{W\varepsilon\hat{H}\left[\frac{1}{r}\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right) + 2\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right)\frac{\partial\ln\phi}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right] + \right. \\ & W\varepsilon\hat{K}\left[v_r\left(\frac{\partial\beta_r\phi^2}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right) - \frac{\alpha}{\phi^2}\frac{\partial\ln\alpha W}{\partial r} + \alpha W^2\left(\beta_r\frac{\partial v_r}{\partial r} - \frac{\partial v_r}{\partial t}\right)\right] - \\ & \left.\varepsilon\hat{L}\left[\frac{\beta_r W}{r} - \frac{\partial\beta_r W}{\partial r} + Wv_{r,r}\frac{\partial}{\partial r}\left(\frac{\alpha}{r\phi^2}\right) + W^3\left(\frac{\alpha}{\phi^2}\frac{\partial v_r}{\partial r} + v_r\frac{\partial v_r}{\partial t}\right)\right]\right\} + \\ & (\hat{J} - \hat{K})\left[v_r\left(\frac{\beta_r}{r} - \frac{\partial\beta_r}{\partial r}\right) + \frac{\partial}{\partial r}\left(\frac{W\alpha}{\phi^2}\right) - \frac{W\alpha}{r\phi^2} + W^3\left(\frac{\partial v_r}{\partial t} - \beta_r\frac{\partial v_r}{\partial r}\right)\right] + \\ & (\hat{H} - \hat{L})\left[\frac{W^3\alpha}{\phi^2}\frac{\partial v_r}{\partial r} + \frac{\beta_r W}{r} - \frac{\partial\beta_r W}{\partial r} - Wv_{r,r}\frac{\partial}{\partial r}\left(\frac{\alpha}{r\phi^2}\right) + \frac{\partial W}{\partial t}\right] - \\ & W\hat{H}\left[\frac{1}{r}\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right) + 2\left(\beta_r - \frac{\alpha v_r}{\phi^2}\right)\frac{\partial\ln\phi}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right] - \\ & W\hat{K}\left[v_r\left(\frac{\partial\beta_r\phi^2}{\partial r} - 2\frac{\partial\ln\phi}{\partial t}\right) - \frac{\alpha}{\phi^2}\frac{\partial\ln\alpha W}{\partial r} + \alpha W^2\left(\beta_r\frac{\partial v_r}{\partial r} - \frac{\partial v_r}{\partial t}\right)\right] + \\ & \hat{L}\left[\frac{\beta_r W}{r} - \frac{\partial\beta_r W}{\partial r} + Wv_{r,r}\frac{\partial}{\partial r}\left(\frac{\alpha}{r\phi^2}\right) + W^3\left(\frac{\alpha}{\phi^2}\frac{\partial v_r}{\partial r} + v_r\frac{\partial v_r}{\partial t}\right)\right] = \alpha\hat{C}^{(1)}. \end{aligned} \quad (2.29)$$

Neutrino Reactions in Supernovae

Beta processes:

- $e^- + p \rightleftharpoons n + \nu_e$
- $e^+ + n \rightleftharpoons p + \bar{\nu}_e$
- $e^- + A \rightleftharpoons \nu_e + A^*$

Neutrino scattering:

- $\nu + n, p \rightleftharpoons \nu + n, p$
- $\nu + A \rightleftharpoons \nu + A$
- $\nu + e^\pm \rightleftharpoons \nu + e^\pm$

Thermal pair processes:

- $N + N \rightleftharpoons N + N + \nu + \bar{\nu}$
- $e^+ + e^- \rightleftharpoons \nu + \bar{\nu}$

Neutrino-neutrino reactions:

- $\nu_x + \nu_e, \bar{\nu}_e \rightleftharpoons \nu_x + \nu_e, \bar{\nu}_e$
($\nu_x = \nu_\mu, \bar{\nu}_\mu, \nu_\tau, \text{ OR } \bar{\nu}_\tau$)
- $\nu_e + \bar{\nu}_e \rightleftharpoons \nu_{\mu,\tau} + \bar{\nu}_{\mu,\tau}$


The Curse and Challenge of the Dimensions

Boltzmann equation determines neutrino distribution function in 6D phase space and time

$$f(r, \theta, \phi, \Theta, \Phi, \epsilon, t)$$

Integration over 3D momentum space yields source terms for hydrodynamics

$$Q(r, \theta, \phi, t), \dot{Y}_e(r, \theta, \phi, t)$$




Solution approach

- **3D** hydro + **6D** direct discretization of Boltzmann Eq. (code development by Sumiyoshi & Yamada '12)
- **3D** hydro + two-moment closure of Boltzmann Eq. (next feasible step to full 3D; cf. Kuroda et al. 2012)
- **3D** hydro + "**ray-by-ray-plus**" variable Eddington factor method (method used at MPA/Garching)
- **2D** hydro + "**ray-by-ray-plus**" variable Eddington factor method (method used at MPA/Garching)

Required resources

- $\geq 10\text{--}100$ PFlops/s (sustained!)
- $\geq 1\text{--}10$ Pflops/s, TBytes
- $\geq 0.1\text{--}1$ PFlops/s, Tbytes
- $\geq 0.1\text{--}1$ Tflops/s, < 1 TByte

"Ray-by-Ray" Approximation for Neutrino Transport in 2D and 3D Geometry



Solve large number of **spherical transport problems** on **radial "rays"** associated with angular zones of polar coordinate grid


Suggests efficient parallelization over the "rays"

radial "ray" →


Performance and Portability of our Supernova Code *Prometheus-Vertex*

- Code employs **hybrid MPI/OpenMP** programming model (collaborative development with **Katharina Benkert, HLRS**).
- Code has been **ported** to different computer platforms by **Andreas Marek, High Level Application Support, Rechenzentrum Garching (RZG)**.
- Code shows **excellent parallel efficiency**, which will be fully exploited in 3D.

Strong Scaling



Relativistic 2D CCSN Explosion Models





Bernhard Müller, THJ, et al. (ApJ 756, ApJ 761, arXiv:1210.6984)

Basic confirmation of previous explosion models for 11.2 and 15 M_{SUN} stars by Marek & THJ (2009)

Relativistic 2D CCSN Explosion Models

"Diagnostic energy" of explosion





Maximum shock radius

Growing set of 2D CCSN Explosion Models



Average shock radius

Florian Hanke (PhD project)





Support for 2D CCSN Explosion Models

2D explosions for 13 M_{sun} progenitor of Nomoto & Hashimoto (1988)



2D explosions for 11.2 and 15 M_{sun} progenitors of Woosley et al. (2002, 1995)





Support for 2D CCSN Explosion Models

AXISYMMETRIC *AB INITIO* CORE-COLLAPSE SUPERNOVA SIMULATIONS OF 12–25 M_{\odot} STARS

STEPHEN W. BRUENN¹, ANTHONY MEZZACAPPA^{2,3,4}, W. RAPHAEL HIX^{2,3}, ERIC J. LENTZ^{3,2,5}, O. E. BRONSON MESSER^{6,3,4}, ERIC J. LINGERFELT^{2,4}, JOHN M. BLONDIN⁷, EIRIK ENDEVE⁴, PEDRO MARRONETTI^{1,8}, AND KONSTANTIN N. YAKUNIN¹

2D explosions for 12, 15, 20, 25 M_{sun} progenitors of Woosley & Heger (2007)



Bruenn et al., arXiv:1212.1747

2D SN Explosion Models



- Basic confirmation of the neutrino-driven mechanism
- Confirm reduction of the critical neutrino luminosity that enables an explosion in self-consistent 2D treatments compared to 1D

However, many aspects are different:

- Different explosion behavior and different explosion energies
- Different codes, neutrino transport and reactions, EoS treatment
- Partly different progenitor sets

Comparisons are urgently needed!


Comparison of 1D & 2D Explosion Models







1D and 2D simulations for 12 M_{sun} progenitor of Woosley & Heger (2007) by F. Hanke (Newtonian with GR gravity corrections) and by B. Müller (GR) agree well with each other.

But Garching models disagree with Bruenn et al. (2012) results!

Garching models are not close to explosion.



Comparison of 1D & 2D Explosion Models



2D SN Explosion Models

Many aspects are different:

- **Different explosion behavior and different explosion energies**
- **Different codes, neutrino transport and reactions, EoS treatment**

Direct comparisons are urgently needed!

Challenge and Goal: 3D

- 2D explosions seem to be “marginal”, at least for some progenitor models and in some (the most?) sophisticated simulations.
- Nature is three dimensional, but 2D models impose the constraint of axisymmetry on the flow!
- Turbulent cascade in 3D transports energy from large to small scales, which is opposite to 2D.
- Is 3D turbulence more supportive to an explosion?
Is the third dimension the key to the neutrino mechanism?
- 3D models are needed to confirm explosion mechanism suggested by 2D simulations!



3D vs. 2D Differences:
The Dimension Conundrum

2D-3D Differences in Parametric Explosion Models

- Nordhaus et al. (ApJ 720 (2010) 694) performed 2D & 3D simulations with **simple neutrino-heating and cooling terms** (no neutrino transport but lightbulb) and found 15–25% improvement in 3D for 15 M_{sun} progenitor star (ApJ 720 (2010) 694)



$$\mathcal{H} = 1.544 \times 10^{20} \left(\frac{L_{\nu_e}}{10^{52} \text{ erg s}^{-1}} \right) \left(\frac{T_{\nu_e}}{4 \text{ MeV}} \right)^2 \times \left(\frac{100 \text{ km}}{r} \right)^2 (Y_n + Y_p) e^{-\tau_{\nu_e}} \left[\frac{\text{erg}}{\text{g s}} \right]$$

$$\mathcal{C} = 1.399 \times 10^{20} \left(\frac{T}{2 \text{ MeV}} \right)^6 (Y_n + Y_p) e^{-\tau_{\nu_e}} \left[\frac{\text{erg}}{\text{g s}} \right]$$



2D-3D Differences in Parametric Explosion Models

- F. Hanke (Diploma Thesis, MPA, 2010) in agreement with L. Scheck (PhD Thesis, MPA, 2007) **could not confirm the findings by Nordhaus et al. (2010)** ! 2D and 3D simulations for $11.2 M_{\text{sun}}$ and $15 M_{\text{sun}}$ progenitors are very similar but results depend on numerical grid resolution: 2D with higher resolution explodes easier, 3D shows opposite trend!




Hanke et al., ApJ 755 (2012) 138,
arXiv:1108.4355


2D & 3D slices for $11.2 M_{\text{sun}}$ model, $L = 1.0 \cdot 10^{52} \text{ erg/s}$

2D-3D Differences

Average entropy of gas in gain layer is not good diagnostic quantity for proximity to explosion



Nordhaus et al., ApJ 720 (2010) 694



Hanke et al., ApJ 755 (2012) 138,
arXiv:1108.4355

Growing "Diversity" of 3D Results

- Dolence et al. (arXiv:1210.5241) find much smaller 2D/3D difference of critical luminosity, but still slightly earlier explosion in 3D.
- Takiwaki et al. (ApJ 749:98, 2012) obtain explosion for an 11.2 M_{sun} progenitor in 3D later than in 2D. Find a bit faster 3D explosion with higher resolution.
- Couch (arXiv:1212.0010) finds also later explosions in 3D than in 2D and higher critical luminosity in 3D!
But critical luminosity increases in 2D with better resolution.
- Ott et al. (arXiv:1210.6674) reject relevance of SASI in 3D and conclude that neutrino-driven convection dominates evolution.

Reasons for 2D/3D differences and different results by different groups are not understood!

Growing "Diversity" of 3D Results

- These results **do not yield a clear picture of 3D effects.**

But:

- The simulations were performed with **different grids** (cartesian+AMR, polar), different codes (CASTRO, ZEUS, FLASH, Cactus, Prometheus), and **different treatments of input physics** for EOS and neutrinos, some **with simplified, not fully self-consistent set-ups.**
- **Resolution differences** are difficult to assess and are likely to strongly depend on spatial region and coordinate direction.
- **Partially compensating effects of opposite influence** might be responsible for the seemingly conflicting results.
- **Convergence tests with much higher resolution** and **detailed code comparisons** for “clean”, well defined problems are **urgently needed**, but both will be ambitious!

Full-Scale 3D Core-Collapse
Supernova Models with Detailed
Neutrino Transport

3D Supernova Models

PRACE grant of 146.7 million core hours allows us to do the first 3D simulations on 16.000 cores.



SuperMUC Petascale System

TGCC Curie



created by LRZ (2012)

Computing Requirements for 2D & 3D Supernova Modeling

Time-dependent simulations: $t \sim 1$ second, $\sim 10^6$ time steps!

CPU-time requirements for one model run:

★ In 2D with 600 radial zones, 1 degree lateral resolution:

$\sim 3 \cdot 10^{18}$ Flops, need $\sim 10^6$ processor-core hours.

★ In 3D with 600 radial zones, 1.5 degrees angular resolution:

$\sim 3 \cdot 10^{20}$ Flops, need $\sim 10^8$ processor-core hours.







John von Neumann
Institut für Computing



3D Core-Collapse Models

27 M_{sun} progenitor






Florian Hanke,
PhD project

3D Core-Collapse Models

27 M_{sun} progenitor



Shock position (max., min., avg.)



Florian Hanke, PhD project

3D Core-Collapse Models

11.2 M_{sun} progenitor




Florian Hanke, PhD project



3D Core-Collapse Models

11.2 M_{sun} progenitor


Shock position (max., min., avg.)



Florian Hanke, PhD project



Numerical Convergence?



Florian Hanke, PhD project

2D simulations are converged; no difference between 0.7, 1.4, and 2.04 degrees angular resolution.

But: 3D simulations may need more resolution for convergence than in 2D!

Numerical Convergence?

Hanke et al., ApJ 755 (2012) 138; arXiv:1108.4355




Figure 16. Turbulent energy spectra $E(l)$ as functions of the multipole order l for different angular resolution. The spectra are based on a decomposition of the azimuthal velocity v_θ into spherical harmonics at radius $r = 150$ km and 400 ms post-bounce time for $15 M_\odot$ runs with an electron–neutrino luminosity of $L_{\nu_e} = 2.2 \times 10^{52}$ erg s⁻¹. Left: 2D models with different angular resolution (black, different thickness) and, for comparison, the 3D model with the highest employed angular resolution (gray). Right: 3D models with different angular resolution and, for comparison, the 2D model with the highest employed angular resolution (gray). The power-law dependence and direction of the energy and enstrophy cascades (see the text) are indicated by red lines and labels for 2D models in the left panel and 3D models in the right panel. The left vertical, dotted line roughly marks the energy-injection scale, and the right vertical, dotted line denotes the onset of dissipation at high l for the best-displayed resolution.

Turbulent energy cascade in 2D from small to large scales, in 3D from large to small scales! =====> **More than 2 degree resolution needed in 3D!**

Summary

- Modelling of SN explosion mechanism has made considerable progress in 1D and multi-D.
- 2D relativistic models yield explosions for “soft” EoSs. Explosion energy tends to be on low side (except recent models by Bruenn et al., arXiv:1212.1747).
- 3D modeling has only begun. No clear picture of 3D effects yet. **But SASI can dominate (during phases) also in 3D models!**
- 3D SN modeling is extremely challenging and variety of approaches for neutrino transport and hydrodynamics/grid choices will be and need to be used.
- Numerical effects (and artifacts) and resolution dependencies in 2D and 3D models must still be understood.
- Bigger computations on faster computers are indispensable, but higher complexity of highly-coupled multi-component problem will demand special care and quality control.