

One-phonon excitations in ^{92}Zr from electron scattering

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2010 J. Phys.: Conf. Ser. 205 012040

(<http://iopscience.iop.org/1742-6596/205/1/012040>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 130.83.133.251

The article was downloaded on 28/07/2013 at 13:54

Please note that [terms and conditions apply](#).

One-phonon excitations in ^{92}Zr from electron scattering

Abdulahman Scheikh Obeid, O Burda, M Chernykh, A Krugmann,
P von Neumann-Cosel, N Pietralla, I Poltoratska, V Ponomarev, and
C Walz

Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

E-mail: scheikh@ikp.tu-darmstadt.de

Abstract. Low-lying collective vibrational excitations of ^{92}Zr have been investigated with electron scattering at the S-DALINAC. The form factors of isospin polarized one-quadrupole phonon states of ^{92}Zr at the $Z = 40$ proton subshell closure have been measured and the momentum-transfer dependence of the form factors for the one-quadrupole phonon states have been compared to the prediction of the Quasiparticle Phonon Model. The E2 transition strengths of the one-quadrupole phonon states and the E3 transition strength of the one-octupole phonon state have been extracted. A comparison to the data on ^{94}Mo and previous spectroscopic information on mixed-symmetric states (MSSs) of ^{92}Zr will be given.

1. Introduction

Collectivity, isospin symmetry and shell structure are generic features of many-body quantum systems that consist of two components (protons and neutrons) like heavy atomic nuclei. The study of collective valence-shell excitations is a useful way to understand how these features coexist, interplay and compete. The investigation of MSSs is an important source of information on the effective proton-neutron interaction in the valence shell [1]. MSSs have been defined in the framework of the Interacting Bosons Model IBM-2 [2]. In analogy to isospin, the symmetry of a multi-boson wave function formed by N_π proton bosons and N_ν neutron bosons with respect to proton and neutron boson labels is quantified by the F-spin. States with $F < F_{max} = (N_\pi + N_\nu)/2$ have wave functions that contain at least one pair of proton and neutron bosons which is antisymmetric under the exchange of proton and neutron labels. They are called mixed-symmetry states. Those states with $F = F_{max}$ are symmetric under the pairwise exchange of the nucleon labels. They are fully symmetric states. The signature for MSSs are (i) strong $M1$ transition to fully symmetric states with matrix elements of about $1\mu_N$ in size and (ii) weakly-collective $E2$ transition to fully symmetric states.

The prediction of the IBM-2 with respect to a multi-phonon structure of MSSs in vibrational nuclei was well confirmed ten years ago by the observation of large $M1$ transitions between low-energy states of ^{94}Mo [3, 4, 5]. The mixed-symmetric 2^+ states were also investigated with electron scattering experiments at the 130 MeV superconducting electron accelerator S-DALINAC [6]. The evaluation of the measured form factors as a function of the momentum-transfer had supported the one-phonon interpretation of the 2_1^+ , 2_3^+ states. In the neighbouring even-even isotone ^{92}Zr which is formed by $N = 52$ neutrons with two valence neutrons and

$Z = 40$ with no valence protons occupying the $\pi(g_{9/2})$ sub-shell a stronger configurational isospin polarization of the one-phonon states than in ^{94}Mo was expected. This has recently been discussed in the literature [7, 8, 9, 10]. In order to deepen the discussions and to verify this expectation a new electron scattering experiment at the S-DALINAC has been performed on the nucleus ^{92}Zr . A calculation of the momentum-transfer dependence of the form factors for the one-quadrupole phonon states has been done in the framework of the Quasiparticle Phonon Model.

2. Experiment and analysis

2.1. Experimental setup and data

The experiment has been carried out at the Darmstadt superconducting electron linear accelerator S-DALINAC which provides electron beams with energy up to 130 MeV and beam current up to $3 \mu\text{A}$. The $33^\circ - 169^\circ$ spectrometer with momentum acceptance of $\pm 2.1\%$ was used. The spectrometer has in its focal plane a detector system based on four single-strip silicon detectors, each provides 96 strips with thickness of $500 \mu\text{m}$ and a pitch of $650 \mu\text{m}$.

We have used an incident beam energy $E_e = 63 \text{ MeV}$ with beam currents from 0.5 to $1 \mu\text{A}$. A ^{92}Zr target with an enrichment of 94.57% and a thickness of 9.75 mg/cm^2 was used. Data were taken at five different scattering angles $\Theta_e = 69^\circ, 81^\circ, 93^\circ, 117^\circ, 165^\circ$ covering the maximum of the $E2$ form factor. Examples of electron spectra are shown in Fig. 1. The salient peaks correspond to the elastic line, the weakly collective one-phonon 2_1^+ state ($E_x = 0.934 \text{ MeV}$), The 3_1^- state ($E_x = 2.339 \text{ MeV}$) and the one-phonon 2_2^+ state ($E_x = 1.847 \text{ MeV}$).




Figure 1. Two electron scattering spectra of the $^{92}\text{Zr}(e,e')$ reaction at $E_e = 63 \text{ MeV}$ and $\Theta_e = 93^\circ, 117^\circ$.

The spectra taken in the $^{92}\text{Zr}(e,e')$ electron scattering reaction were energy calibrated with the excitation energies of the $0_1^+, 2_1^+, 2_2^+, 3_1^-$ states given in [9].

2.2. Data analysis

The experimentally extracted cross sections for the excited states were normalized to the cross section of the elastic line. The absolute values of the differential cross sections were calculated

from the corresponding peak areas taking into account the radiative corrections and the dead time of the electronics according to

$$\left(\frac{d\sigma}{d\Omega}\right) = A_{in}^{exp} \cdot \frac{1}{\Delta\Omega} \cdot \frac{e}{It} \cdot \frac{\mu_{mol}}{n_{eff}N_A} \quad (1)$$

were $\Delta\Omega$ is the spectrometer solid angle [sr], It the accumulated charge [C], μ_{mol} the molecular mass, [g/mol], n_{eff} the effective areal density of the target [g/cm²] and N_A the Avogadro number [1/mol].

The squared form factor is given as:

$$|F(E_0, \theta)|^2 = \frac{(d\sigma/d\Omega)_{exp}}{(d\sigma/d\Omega)_{Mott}} \quad (2)$$




Figure 2. Momentum-transfer dependence of the form factor of the one-phonon FS 2_1^+ state (top), MS 2_2^+ state (middle) and 3_1^- state (bottom) of ⁹²Zr from electron scattering. The data (full squares) are compared to the QPM (solid lines).

with the Mott cross section:

$$\left(\frac{d\sigma}{d\Omega}\right)_{Mott} = \left(\frac{Ze^2}{2E_0}\right)^2 \frac{\cos^2(\theta/2)}{\sin^4(\theta/2)} \quad (3)$$

Calculations of the momentum-transfer dependence of the form factor of the 2^+ states and of the 3_1^- state have been performed within the Quasiparticle Phonon Model (QPM). Excited states of ^{92}Zr have been described by a wave function which includes one-, two-, and three-phonon configurations. The (e,e') cross sections have been calculated within the Distorted Wave Born Approximation using the code of Heisenberg and Blok [11] and employing the radial transition charge densities calculated by the QPM. The E2 and E3 strengths can be extrapolated to the photon point, $q = E_x/\hbar c$.

Table 1. E2 strengths from (e,e') in comparison to the known values from $(n,n'\gamma)$ [5, 9] and the E3 strength from (e,e') in W.u.

	This work: $^{92}\text{Zr}(e,e')$ E λ strength in W.u. Form factor scaled to QPM	$^{92}\text{Zr}(n,n'\gamma)$ [9]	$^{94}\text{Mo}(n,n'\gamma)$ [5]
$B(E2; 2_1^+ \rightarrow 0_1^+)$	6.2 ± 0.3	6.4 ± 0.5	16.0 ± 0.2
$B(E2; 2_2^+ \rightarrow 0_1^+)$	3.3 ± 0.2	3.4 ± 0.4	2.2 ± 0.2
$B(E3; 3_1^- \rightarrow 0_1^+)$	18.3 ± 1.1		

The $B(E2; 2_1^+ \rightarrow 0_1^+)$ and $B(E2; 2_2^+ \rightarrow 0_1^+)$ values that were extracted using the form factor calculated with the QPM are in good agreement with the literature values [9]. The EM strengths are compared to the literature and to the ones of the symmetric and mixed-symmetric states of ^{94}Mo in Tab. 1. The extracted value of the $B(E2; 3_1^- \rightarrow 0_1^+)$ is 18.1 ± 1.1 W.u. and it compares well to the values obtained from proton scattering experiments (14.7, 16.2, 18.9, 21.3, 23.6) W.u in the review article by Spear [12].

3. Discussion

The (e,e') electron scattering reaction is predominantly a one-step process. Fig. 1 shows that the 2_2^+ state of ^{92}Zr is almost as strongly populated in (e,e') as the one-phonon 2_1^+ state suggesting a large one-phonon component being present in the wave function of the 2_2^+ state either. Figure 2 presents the form factors of these 2^+ states and of the 3_1^- state as a function of the momentum transfer q . The similarity of the momentum-transfer dependence of the form factors between the 2^+ states further supports the one-phonon nature of the 2_2^+ state of ^{92}Zr .

The theoretical results shown in Fig. 2 are absolute and not scaled to the experimental data. The QPM results provide a good description of the (e,e') form factor in this interval of the momentum transfer. The main configurations of the wave functions used to calculate the form factors are given as following:

$$\begin{aligned}\psi_{2_1^+} &= \sqrt{0.59} \nu \left(2d_{5/2}^2 \right) + \sqrt{0.14} \pi \left(1g_{9/2}^2 \right) \\ \psi_{2_2^+} &= -\sqrt{0.38} \nu \left(2d_{5/2}^2 \right) + \sqrt{0.27} \pi \left(1g_{9/2}^2 \right)\end{aligned}\quad (4)$$

The QPM calculations predict the main proton and neutron components in the wave functions of the one-phonon 2^+ states to have the opposite relative signs which confirms the picture of symmetric and mixed-symmetric character in the valance shell. The 2_1^+ state, because of the large neutron component, is a predominantly symmetric albeit isospin polarized state and the 2_2^+ state is a mixed-symmetric one with a weak isospin polarization.

The E2 strengths extracted in this work are close to the values, $6.4_{-0.5}^{+0.6}$ and $3.4(4)$ for the 2_1^+ and 2_2^+ respectively, that were obtained from a QPM calculations performed in Ref. [10]. However,

the wave functions published for the 2^+ states in Ref. [10] show stronger isospin polarisation of the 2_1^+ and 2_2^+ states than what we have found from our calculation.

Acknowledgments

We thank Ch Stoyanov, N Lo Iudice, V Werner, C Fransen, G Rainovski and D Tarpanov for discussions. Support by the DFG under Grant No. SFB 634, by the Helmholtz International Center for FAIR and by the DAAD within the German-Bulgarian exchange program under grant Nos. D/06/05918 and DAAD-09 is gratefully acknowledged.

References

- [1] Pietralla N, von Brentano P and Lisetskiy A F 2008 *Prog. Part. Nucl. Phys.* **60** 225
- [2] Iachello F and Arima A 1987 *The Interacting Boson Model* (Cambridge University Press.)
- [3] Pietralla N *et al.* 1999 *Phys. Rev. Lett.* **83** 1303
- [4] Pietralla N, Fransen C, von Brentano P, Dewald A, Fitzler A, Frießner C and Gableske J 2000 *Phys. Rev. Lett.* **84** 3775
- [5] Fransen C *et al.* 2003 *Phys. Rev C* **67** 024307
- [6] Burda O *et al.* 2007 *Phys. Rev. Lett* **99** 029503
- [7] Werner V *et al.* 2002 *Phys. Lett. A* **550** 140
- [8] Holt J D, Pietralla N, Holt J W, Kuo T T S and Rainovski G 2007 *Phys. Rev. C* **76** 034325
- [9] Fransen C *et al.* 2005 *Phys. Rev C* **71** 054304
- [10] LoIudice N and Stoyanov C 2006 *Phys. Rev. C* **73** 037305
- [11] Heisenberg J and Blok H P 1983 *Annu. Rev. Nucl. Part. Sci. C* **33** 0569
- [12] Spear R H 1989 *Atom. Dat. And Nuc. Dat. Tab* **42** 55