

Ab Initio Calculations of Nuclear Structure

Lecture 1: Hamiltonian

Robert Roth

Institut für Kernphysik - Theoriezentrum



TECHNISCHE
UNIVERSITÄT
DARMSTADT

HFHF Helmholtz
Forschungsakademie
Hessen für FAIR

Overview

■ **Lecture 1: Hamiltonian**

Prelude • Many-Body Quantum Mechanics • Nuclear Hamiltonian • Matrix Elements • Two-Body Problem • Correlations & Unitary Transformations

■ **Lecture 2: Light Nuclei**

Similarity Renormalization Group • Many-Body Problem • Configuration Interaction • No-Core Shell Model • Basis Optimization

■ **Lecture 3: Medium-Mass Nuclei**

Normal Ordering • Coupled-Cluster Theory • In-Medium Similarity Renormalization Group • Many-Body Perturbation Theory

■ **Project: Do-It-Yourself NCSM**

Three-Body Problem • Numerical SRG Evolution • NCSM Eigenvalue Problem • Lanczos Algorithm

■ **Lecture 4: Precision, Uncertainties, and Applications**

Chiral Interactions for Precision Calculations • Uncertainty Quantification • Applications to Nuclei and Hypernuclei


Prelude

Theoretical Context

better resolution / more fundamental

Quantum Chromodynamics


Nuclear Structure



- finite nuclei
- few-nucleon systems
- nuclear interaction
- hadron structure
- quarks & gluons
- deconfinement

The Problem

$$H |\Psi_n\rangle = E_n |\Psi_n\rangle$$



The Problem

$$H |\Psi_n\rangle = E_n |\Psi_n\rangle$$

ab initio :=

solve nuclear many-body problem
based on realistic interactions
using controlled and improvable truncations
with quantified theoretical uncertainties

The Problem

$$H |\Psi_n\rangle = E_n |\Psi_n\rangle$$

Assumptions

- use nucleons as effective degrees of freedom
- use non-relativistic framework, relativistic corrections are absorbed in Hamiltonian
- use Hamiltonian formulation, i.e., conventional many-body quantum mechanics
- focus on bound states, though continuum aspects are very interesting

The Problem

$$H |\Psi_n\rangle = E_n |\Psi_n\rangle$$

What is this many-body Hamiltonian?

nuclear forces, chiral effective field theory, three-body interactions, consistency, convergence,...

What about these many-body states?

many-body quantum mechanics, antisymmetry, second quantisation, many-body basis, truncations,...

How to solve this equation?

ab initio methods, correlations, similarity transformations, large-scale diagonalization, coupled-cluster theory,...

Many-Body Quantum Mechanics

... a very quick reminder

Single-Particle Basis

- effective constituents are nucleons characterized by **position, spin and isospin** degrees of freedom

$$|\alpha\rangle = |\text{position}\rangle \otimes |\text{spin}\rangle \otimes |\text{isospin}\rangle$$

- typical **basis choice** for configuration-type bound-state methods

$|\text{position}\rangle = |nlm_l\rangle$ spherical harmonic oscillator or other spherical single-particle potential

$|\text{spin}\rangle = |s = \frac{1}{2}, m_s\rangle$ eigenstates of s^2 and s_z with $s=1/2$

$|\text{isospin}\rangle = |t = \frac{1}{2}, m_t\rangle$ eigenstates of t^2 and t_3 with $t=1/2$

- use **spin-orbit coupling** at the single-particle level

$$|n(l\frac{1}{2})jm; \frac{1}{2}m_t\rangle = \sum_{m_l, m_s} c \left(\begin{array}{cc|c} l & 1/2 & j \\ m_l & m_s & m \end{array} \right) |nlm_l\rangle \otimes |\frac{1}{2}m_s\rangle \otimes |\frac{1}{2}m_t\rangle$$

Identical Particles & Spin-Statistics Theorem

- **systems of identical particles**: many-body states have to be eigenstates of the transposition operator for any particle pair with eigenvalues ± 1

$$T_{ij} |\psi\rangle = +1 |\psi\rangle$$

states symmetric under transposition of any pair of particle indices

$$T_{ij} |\psi\rangle = -1 |\psi\rangle$$

states antisymmetric under transposition of any pair of particles

- simple **product states** are not suitable for systems of identical particles

$$|\phi\rangle = |\alpha_1\rangle \otimes |\alpha_2\rangle \otimes \cdots \otimes |\alpha_A\rangle$$

- **spin-statistics theorem** connects transposition symmetry to particle spin:
 - bosons = integer spin = symmetric states
 - fermions = half-integer spin = antisymmetric states

- focus on fermions, i.e., **antisymmetric states in the following**

Slater Determinants

- antisymmetric states can be constructed via the **antisymmetrization operator**

$$\mathcal{A} = \frac{1}{A!} \sum_{\pi} \text{sgn}(\pi) P_{\pi}$$

sum over all permutations signum of permutation permutation operator

- technically it is a **projection operator** onto the antisymmetric A-body Hilbert space and has the same structure as a **general determinant**
- **Slater determinants**: antisymmetrized product states

$$\begin{aligned} |\alpha_1 \alpha_2 \dots \alpha_A\rangle &= \sqrt{A!} \mathcal{A} (|\alpha_1\rangle \otimes |\alpha_2\rangle \otimes \dots \otimes |\alpha_A\rangle) \\ &= \frac{1}{\sqrt{A!}} \sum_{\pi} \text{sgn}(\pi) P_{\pi} (|\alpha_1\rangle \otimes |\alpha_2\rangle \otimes \dots \otimes |\alpha_A\rangle) \end{aligned}$$

- **Pauli principle is a consequence of antisymmetry**: you cannot antisymmetrize a product state that contains two identical single-particle states

Slater Determinants as Basis

- given a complete single-particle basis $\{|\alpha\rangle\}$ then the set of Slater determinants formed by all possible combinations of A different single-particle states is a **complete basis of the antisymmetric A-body Hilbert space**

- resolution of the **identity operator**

$$1 = \sum_{\alpha_1 < \alpha_2 < \dots < \alpha_A} |\alpha_1 \alpha_2 \dots \alpha_A\rangle \langle \alpha_1 \alpha_2 \dots \alpha_A| = \frac{1}{A!} \sum_{\alpha_1, \alpha_2, \dots, \alpha_A} |\alpha_1 \alpha_2 \dots \alpha_A\rangle \langle \alpha_1 \alpha_2 \dots \alpha_A|$$

- careful with **double counting**: Slater determinants that differ only by the order of the single-particle states are identical up to a sign...

- **expansion of general antisymmetric state** in Slater determinant basis

$$|\Psi\rangle = \sum_{\alpha_1 < \alpha_2 < \dots < \alpha_A} C_{\alpha_1 \alpha_2 \dots \alpha_A} |\alpha_1 \alpha_2 \dots \alpha_A\rangle = \sum_i C_i |\{\alpha_1 \alpha_2 \dots \alpha_A\}_i\rangle$$

Second Quantization: Basics

- define **Fock-space** as direct sum of A-particle Hilbert spaces

$$\mathcal{F} = \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots \oplus \mathcal{H}_A \oplus \cdots$$

- **vacuum state**: the only state in the zero-particle Hilbert space

$$|0\rangle \in \mathcal{H}_0 \quad \langle 0|0\rangle = 1 \quad |0\rangle \neq 0$$

- **creation operators**: add a particle in single-particle state $|\alpha\rangle$ to an A-body Slater determinant yielding an (A+1)-body Slater determinant

$$a_{\alpha}^{\dagger} |0\rangle = |\alpha\rangle$$

$$a_{\alpha}^{\dagger} |\alpha_1 \alpha_2 \dots \alpha_A\rangle = \begin{cases} |\alpha \alpha_1 \alpha_2 \dots \alpha_A\rangle & ; \alpha \notin \{\alpha_1 \alpha_2 \dots \alpha_A\} \\ 0 & ; \text{otherwise} \end{cases}$$

- resulting states are automatically normalized and antisymmetrized
- new single-particle state is added in the first slot, can be moved elsewhere through transpositions

Second Quantization: Basics

- **annihilation operators**: remove a particle with single-particle state $|\alpha\rangle$ from an A-body Slater determinant yielding an (A-1)-body Slater determinant

$$a_\alpha |0\rangle = 0$$

$$a_\alpha |\alpha_1 \alpha_2 \dots \alpha_A\rangle = \begin{cases} (-1)^{i-1} |\alpha_1 \alpha_2 \dots \alpha_{i-1} \alpha_{i+1} \dots \alpha_A\rangle & ; \alpha = \alpha_i \\ 0 & ; \text{otherwise} \end{cases}$$

- annihilation operator acts on first slot, need transpositions to get correct single-particle state there
- based on these definitions one can easily show that creation and annihilations operators satisfy **anticommutation relations**

$$\{a_\alpha, a_{\alpha'}\} = 0$$

$$\{a_\alpha^\dagger, a_{\alpha'}^\dagger\} = 0$$

$$\{a_\alpha, a_{\alpha'}^\dagger\} = \delta_{\alpha\alpha'}$$

- complication of handling permutations in "first quantization" are translated to the commutation behaviour of strings of operators

Second Quantization: States

- Slater determinants can be written as **string of creation operators** acting on vacuum state

$$|\alpha_1 \alpha_2 \dots \alpha_A\rangle = a_{\alpha_1}^\dagger a_{\alpha_2}^\dagger \dots a_{\alpha_A}^\dagger |0\rangle$$

- alternatively one can define an A-body **reference Slater determinant**

$$|\Phi\rangle = |\alpha_1 \alpha_2 \dots \alpha_A\rangle = a_{\alpha_1}^\dagger a_{\alpha_2}^\dagger \dots a_{\alpha_A}^\dagger |0\rangle$$

and construct arbitrary Slater determinants through **particle-hole excitations** on top of the reference state

$$\begin{aligned} |\Phi_a^p\rangle &= a_{\alpha_p}^\dagger a_{\alpha_a} |\Phi\rangle \\ |\Phi_{ab}^{pq}\rangle &= a_{\alpha_p}^\dagger a_{\alpha_q}^\dagger a_{\alpha_b} a_{\alpha_a} |\Phi\rangle \\ &\vdots \end{aligned}$$

index convention: a, b, c, \dots : hole states, occupied in reference state
 p, q, r, \dots : particle state, unoccupied in reference states

Second Quantization: Operators

- **operators** can be expressed in terms of creation and annihilation operators as well, e.g., for one-body kinetic energy and two-body interactions:

'first quantization'

$$T = \sum_{i=1}^A t_i$$

$$V = \sum_{i < j=1}^A v_{ij}$$

second quantization

$$T = \sum_{\alpha\alpha'} \langle \alpha | t | \alpha' \rangle a_{\alpha}^{\dagger} a_{\alpha'}$$

$$V = \frac{1}{4} \sum_{\alpha_1 \alpha_2 \alpha'_1 \alpha'_2} \langle \alpha_1 \alpha_2 | v | \alpha'_1 \alpha'_2 \rangle a_{\alpha_1}^{\dagger} a_{\alpha_2}^{\dagger} a_{\alpha'_2} a_{\alpha'_1}$$

- **set of one or two-body matrix elements** fully defines the one or two-body operator in Fock space
- second quantization is extremely convenient to **compute matrix elements** of operators with Slater determinants

Nuclear Hamiltonian

Nuclear Hamiltonian

- general form of **many-body Hamiltonian** can be split into a center-of-mass and an intrinsic part

$$\begin{aligned} H &= T + V_{NN} + V_{3N} + \dots = T_{\text{cm}} + T_{\text{int}} + V_{NN} + V_{3N} + \dots \\ &= T_{\text{cm}} + H_{\text{int}} \end{aligned}$$


- **intrinsic Hamiltonian** is invariant under translation, rotation, Galilei boost, parity, time evolution, time reversal,...

$$\begin{aligned} H_{\text{int}} &= T_{\text{int}} + V_{NN} + V_{3N} + \dots \\ &= \sum_{i<j}^A \frac{1}{2mA} (\vec{p}_i - \vec{p}_j)^2 + \sum_{i<j}^A v_{NN,ij} + \sum_{i<j<k}^A v_{3N,ijk} + \dots \end{aligned}$$

- these symmetries constrain the possible operator structures that can appear in the interaction terms...

... but how can we really **determine the nuclear interaction** ?

Nature of the Nuclear Interaction



~ 1.6fm

$\sim 1.6\text{fm}$


$$\rho_0^{-1/3} = 1.8\text{fm}$$

- nuclear interaction is **not fundamental**
- residual force analogous to **van der Waals interaction** between neutral atoms
- **based on QCD** and induced via polarization of quark and gluon distributions of nucleons
- **encapsulates all the complications** of the QCD dynamics and the structure of nucleons
- acts only if the nucleons overlap, i.e. at **short ranges**
- irreducible **three-nucleon interactions** are important

Yesterday... from Phenomenology


Wiringa, Machleidt,...

- until 2005: **high-precision phenomenological NN interactions** were state-of-the-art in ab initio nuclear structure theory
 - **Argonne V18**: long-range one-pion exchange plus phenomenological parametrization of medium- and short-range terms, local operator form
 - **CD Bonn 2000**: more systematic one meson-exchange parametrization including pseudo-scalar, scalar and vector mesons, inherently nonlocal
- parameters of the NN potential (~ 40) **fit to NN phase shifts** up to ~ 300 MeV and reproduce them with high accuracy
- supplemented by **phenomenological 3N interactions** consisting of a Fujita-Miyazawa-type term plus various hand-picked contributions
- **fit to ground states and spectra of light nuclei**, sometimes up to $A \leq 8$




Argonne V18 Potential

Wiringa, et al., PRC 51, 38 (1995)




$$\begin{aligned}
 v_{NN} = & \sum_{S,T} v_{ST}^c(r) \Pi_{ST} + \sum_{S,T} v_{ST}^{l^2}(r) \vec{L}^2 \Pi_{ST} \\
 & + \sum_T v_T^t(r) S_{12} \Pi_{1T} + \sum_T v_T^{ls}(r) (\vec{L} \cdot \vec{S}) \Pi_{1T} \\
 & + \sum_T v_T^{ls^2}(r) (\vec{L} \cdot \vec{S})^2 \Pi_{1T} + \dots
 \end{aligned}$$



Tomorrow... from Lattice QCD

Hatsuda, Aoki, Ishii, Beane, Savage, Bedaque,...




N. Ishii et al., PRL 99, 022001 (2007)

- first attempts towards construction of nuclear interactions directly from **lattice QCD simulations**
- compute relative **two-nucleon wave function** on the lattice
- **invert Schrödinger equation** to extract effective two-nucleon potential
- only **schematic results** so far (unphysical masses and mass dependence, model dependence,...)
- **alternatives**: phase-shifts or low-energy constants from lattice QCD

Today... from Chiral EFT

Weinberg, van Kolck, Machleidt, Entem, Meißner, Epelbaum, Krebs, Bernard,...

- low-energy **effective field theory** for relevant degrees of freedom (π, N) based on symmetries of QCD
- explicit long-range **pion dynamics**
- unresolved short-range physics absorbed in **contact terms**, low-energy constants fit to experiment
- systematic expansion in a small parameter with power counting enable **controlled improvements** and **error quantification**
- hierarchy of **consistent NN, 3N, 4N,...** interactions
- consistent **electromagnetic and weak operators** can be constructed in the same framework



Many Choices...

■ **standard chiral NN+3N**

- NN: N3LO, Entem&Machleidt, nonlocal, cutoff 500 MeV
- 3N: N2LO, Navratil, local, cutoff 500 (400) MeV

first generation, most widely used up to now

■ **N2LO-opt, N2LO-sat, ...**

- NN: N2LO, Ekström+, nonlocal, cutoff 500 MeV
- 3N: N2LO, Ekström+, nonlocal, cutoff 500 MeV

improved fitting, also many-body inputs

■ **local N2LO**

- NN: N2LO, Gezerlis+, local, cutoff 1.0...1.2 fm
- 3N: N2LO, Gezerlis+, local, cutoff 1.0...1.2 fm

designed specifically for QMC applications

■ **nonlocal LO...N4LO**

- NN: LO...N4LO, Machleidt, nonlocal, cutoff 450...550 MeV
- 3N: N2LO...N3LO, Hüther+, nonlocal, cutoff 450...550 MeV

the future...

■ **semilocal LO...N4LO+**

- NN: LO...N4LO, Epelbaum, semilocal, cutoff 0.8...1.2 fm
- 3N: N2LO..., LENPIC, semilocal, cutoff 0.8...1.2 fm


...systematic variation of chiral order enables quantification of theory uncertainties

Momentum-Space Matrix Elements


$$\langle q(LS)JM; TM_T | v_{NN} | q'(L'S)JM; TM_T \rangle$$

Argonne V18



J=1
L=0
L'=0
S=1
T=0



chiral NN
(N3LO, E&M, 500 MeV)



J=1
L=0
L'=2
S=1
T=0



Matrix Elements

Partial-Wave Matrix Elements

- **relative partial-wave matrix elements** of NN and 3N interaction are **universal input** for many-body calculations
- selection of **relevant partial-wave bases** in two and three-body space with all M quantum numbers suppressed:

two-body relative momentum: $|q (LS) JT\rangle$

two-body relative HO: $|N (LS) JT\rangle$

three-body Jacobi momentum: $|\pi_1 \pi_2; [(L_1 S_1) J_1, (L_2 \frac{1}{2}) J_2] J_{12}; (T_1 \frac{1}{2}) T_{12}\rangle$

three-body Jacobi HO: $|N_1 N_2; [(L_1 S_1) J_1, (L_2 \frac{1}{2}) J_2] J_{12}; (T_1 \frac{1}{2}) T_{12}\rangle$

antisym. three-body Jacobi HO: $|E_{12} i J_{12}^\pi T_{12}\rangle$

- lots of **transformations** between the different bases are needed in practice
- **exception**: Quantum Monte Carlo methods working in coordinate representation need local operator form

Symmetries and Matrix Elements

- relative partial-wave matrix elements make **maximum use of the symmetries** of the nuclear interaction
- consider, e.g., the relative two-body matrix elements in HO basis

$$\langle N (LS) JM; TM_T | v_{NN} | N' (L'S') J'M'; T'M'_T \rangle$$

- the matrix elements of the NN interaction
 - ... do not connect different J
 - ... do not connect different M and are independent of M
 - ... do not connect different parities
 - ... do not connect different S
 - ... do not connect different T
 - ... do not connect different M_T

$$\Rightarrow \langle N (LS) J; TM_T | v_{NN} | N' (L'S) J; TM_T \rangle$$

- relative matrix elements are **efficient and simple to compute**

Transformation to Single-Particle Basis

- most many-body calculations need **matrix elements with single-particle quantum numbers** (cf. second quantization)

$$\begin{aligned} \langle \alpha_1 \alpha_2 | v_{NN} | \alpha'_1 \alpha'_2 \rangle &= \\ &= \langle n_1 l_1 j_1 m_1 m_{t1}, n_2 l_2 j_2 m_2 m_{t2} | v_{NN} | n'_1 l'_1 j'_1 m'_1 m'_{t1}, n'_2 l'_2 j'_2 m'_2 m'_{t2} \rangle \end{aligned}$$

- obtained from relative HO matrix elements via **Moshinsky-transformation**

$$\begin{aligned} \langle n_1 l_1 j_1, n_2 l_2 j_2; JT | v_{NN} | n'_1 l'_1 j'_1, n'_2 l'_2 j'_2; JT \rangle &= \\ &= \sqrt{(2j_1 + 1)(2j_2 + 1)(2j'_1 + 1)(2j'_2 + 1)} \sum_{\lambda, S, T} \sum_{\lambda', S', T'} \langle \lambda S T | v_{NN} | \lambda' S' T' \rangle \\ &\times \langle l_1 l_2 j_1 j_2 | \lambda S T \rangle \langle l'_1 l'_2 j'_1 j'_2 | \lambda' S' T' \rangle \\ &\times \langle N \Lambda, \nu \lambda | n'_1 l'_1, n'_2 l'_2; L' \rangle \\ &\times (2j_1 + 1)(2S + 1)(2L + 1)(2L' + 1) (-1)^{L+L'} \{1 - (-1)^{\lambda+S+T}\} \\ &\times \langle \nu(\lambda S) j T | v_{NN} | \nu'(\lambda' S') j T \rangle \end{aligned}$$

this analytic transformation from relative to single-particle matrix elements only exists for the harmonic oscillator basis

Matrix Element Machinery

- beneath any ab initio many-body method there is a **machinery for computing, transforming and storing matrix elements** of all operators entering the calculation

compute and store relative two-body HO matrix elements of NN interaction

compute and store Jacobi three-body HO matrix elements of 3N interaction

perform unitary transformations of the two- and three-body relative matrix elements (e.g. Similarity Renormalization Group)

transform to single-particle JT-coupled two-body HO matrix elements and store

transform to single-particle JT-coupled three-body HO matrix elements and store

● ● ●
same for 4N with four-body matrix elements

Two-Body Problem

Solving the Two-Body Problem

- **simplest ab initio problem**: the only two-nucleon bound state, the deuteron
- start from **Hamiltonian in two-body space**, change to center of mass and intrinsic coordinates

$$\begin{aligned} H &= H_{\text{cm}} + H_{\text{int}} = T_{\text{cm}} + T_{\text{int}} + V_{\text{NN}} \\ &= \frac{1}{2M} \vec{p}_{\text{cm}}^2 + \frac{1}{2\mu} \vec{q}^2 + V_{\text{NN}} \end{aligned}$$

- **separate** two-body state into center of mass and intrinsic part

$$|\psi\rangle = |\Phi_{\text{cm}}\rangle \otimes |\phi_{\text{int}}\rangle$$

- solve **eigenvalue problem for intrinsic part** (effective one-body problem)

$$H_{\text{int}} |\phi_{\text{int}}\rangle = E |\phi_{\text{int}}\rangle$$

Solving the Two-Body Problem

- expand eigenstates in a **relative partial-wave HO basis**

$$|\phi_{\text{int}}\rangle = \sum_{NLSJMTM_T} C_{NLSJMTM_T} |N(LS)JM; TM_T\rangle$$

$$|N(LS)JM; TM_T\rangle = \sum_{M_L M_S} c \left(\begin{matrix} L & S \\ M_L & M_S \end{matrix} \middle| \begin{matrix} J \\ M \end{matrix} \right) |NLM_L\rangle \otimes |SM_S\rangle \otimes |TM_T\rangle$$

- **symmetries** simplify the problem dramatically:
 - H_{int} does not connect/mix different J, M, S, T, M_T and parity π
 - angular mom. coupling only allows $J=L+1, L, L-1$ for $S=1$ or $J=L$ for $S=0$
 - total antisymmetry requires $L+S+T=\text{odd}$
- for given J^n at most two sets of angular-spin-isospin quantum numbers contribute to the expansion

Deuteron Problem

- assume $J^\pi = 1^+$ for the **deuteron ground state**, then the basis expansion reduces to

$$|\phi_{\text{int}}, J^\pi = 1^+\rangle = \sum_N C_N^{(0)} |N(01) 1M; 00\rangle + \sum_N C_N^{(2)} |N(21) 1M; 00\rangle$$

- inserting into Schrödinger equation and multiplying with basis bra leads to **matrix eigenvalue problem**


$$\begin{pmatrix} \langle N'(01)\dots | H_{\text{int}} | N(01)\dots \rangle & \langle N'(01)\dots | H_{\text{int}} | N(21)\dots \rangle \\ \langle N'(21)\dots | H_{\text{int}} | N(01)\dots \rangle & \langle N'(21)\dots | H_{\text{int}} | N(21)\dots \rangle \end{pmatrix} \begin{pmatrix} C_{N'}^{(0)} \\ C_{N'}^{(2)} \end{pmatrix} = E \begin{pmatrix} C_{N'}^{(0)} \\ C_{N'}^{(2)} \end{pmatrix}$$

simplest possible Jacobi-NCSM calculation


- eigenvectors ψ coefficients and eigenvalues the energies
- **truncate** N **to** $N \leq N_{\text{max}}$ and choose N_{max} large enough so that observables are converged, i.e., do not depend on N_{max} anymore

Deuteron Solution

Argonne V18



chiral NN



- deuteron wave function show two characteristics that are **signatures of correlations** in the two-body system:
 - suppression at small distances due to short-range repulsion
 - $L=2$ admixture generated by tensor part of the NN interaction

Correlations & Unitary Transformations

Correlations

**correlations:
everything beyond the independent
particle picture**

- many-body eigenstates of independent-particle models described by one-body Hamiltonians are **Slater determinants**
- thus, a single Slater determinant **does not describe correlations**
- but Slater determinants are a basis of the antisym. A-body Hilbert space, so any state can be expanded in Slater determinants
- to describe **short-range correlations**, a superposition of many Slater determinants is necessary

Why Unitary Transformations ?

realistic nuclear interactions generate strong short-range correlations in many-body states

Unitary Transformations

- adapt Hamiltonian to truncated low-energy model space
- improve convergence of many-body calculations
- preserve the physics of the initial Hamiltonian and all observables

many-body methods rely on truncated Hilbert spaces
not capable of describing these correlations

Unitary Transformations

- unitary transformations **conserve the spectrum** of the Hamiltonian, with a unitary operator U we get

$$\begin{aligned} H|\psi\rangle &= E|\psi\rangle & 1 &= U^\dagger U = U U^\dagger \\ U^\dagger H U U^\dagger |\psi\rangle &= E U^\dagger |\psi\rangle & \text{with} & \tilde{H} = U^\dagger H U \\ \tilde{H}|\tilde{\psi}\rangle &= E|\tilde{\psi}\rangle & & |\tilde{\psi}\rangle = U^\dagger |\psi\rangle \end{aligned}$$

- for **other observables** defined via matrix elements of an operator A with the eigenstates we obtain

$$\langle\psi|A|\psi'\rangle = \langle\psi|U U^\dagger A U U^\dagger |\psi'\rangle = \langle\tilde{\psi}|\tilde{A}|\tilde{\psi}'\rangle$$

unitary transformations conserve all observables as long as the Hamiltonian and all other operators are transformed consistently

Overview

■ **Lecture 1: Hamiltonian**

Prelude • Many-Body Quantum Mechanics • Nuclear Hamiltonian • Matrix Elements • Two-Body Problem • Correlations & Unitary Transformations

■ **Lecture 2: Light Nuclei**

Similarity Renormalization Group • Many-Body Problem • Configuration Interaction • No-Core Shell Model • Basis Optimization

■ **Lecture 3: Medium-Mass Nuclei**

Normal Ordering • Coupled-Cluster Theory • In-Medium Similarity Renormalization Group • Many-Body Perturbation Theory

■ **Project: Do-It-Yourself NCSM**

Three-Body Problem • Numerical SRG Evolution • NCSM Eigenvalue Problem • Lanczos Algorithm

■ **Lecture 4: Precision, Uncertainties, and Applications**

Chiral Interactions for Precision Calculations • Uncertainty Quantification • Applications to Nuclei and Hypernuclei