

# Modern Nuclear Structure Theory

From Realistic Interactions to the Nuclear Chart



Robert Roth

Institut für Kernphysik  
Technische Universität Darmstadt

# Overview

## ■ Motivation


## ■ Modern Effective Interactions

- Correlations & Unitary Correlation Operator Method


## ■ Applications

- No Core Shell Model
- Hartree-Fock & Beyond
- Random Phase Approximation & Beyond
- Fermionic Molecular Dynamics

# Nuclear Structure in the 21<sup>st</sup> Century



# Modern Nuclear Structure Theory



# Realistic NN-Potentials

## ■ QCD motivated

- symmetries, meson-exchange picture
- chiral effective field theory

Argonne V18

## ■ short-range phenomenology

- short-range parametrisation or contact terms

CD Bonn

Nijmegen I/II

Chiral N3LO

## ■ experimental two-body data

- scattering phase-shifts & deuteron properties reproduced with high precision


Argonne V18 +  
Illinois 2

## ■ supplementary three-nucleon force

- adjusted to spectra of light nuclei


Chiral N3LO +  
N2LO

# Argonne V18 Potential




# Deuteron: Manifestation of Correlations

$$M_S = 0 \\ \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$$



$$M_S = \pm 1 \\ |\uparrow\uparrow\rangle, |\downarrow\downarrow\rangle$$



- spin-projected two-body density  $\rho_{1,M_S}^{(2)}(\vec{r})$
- **exact deuteron solution** for Argonne V18 potential


two-body density fully suppressed at small particle distances  $|\vec{r}|$

**central correlations**


angular distribution depends strongly on relative spin orientation

**tensor correlations**


# *Ab initio* Methods: GFMC



# Modern Nuclear Structure Theory



# Modern Nuclear Structure Theory



# Why Effective Interactions?

## Realistic Potentials

- generate strong correlations in many-body states
- short-range central & tensor correlations most important

## Many-Body Methods

- rely on truncated many-nucleon Hilbert spaces for  $A > 12$
- not capable of describing short-range correlations
- extreme: Hartree-Fock based on single Slater determinant

## Modern Effective Interactions

- adapt realistic potential to the available model space
- conserve experimentally constrained properties (phase shifts)



# Unitary Correlation Operator Method (UCOM)

# Unitary Correlation Operator Method

## Correlation Operator

introduce short-range correlations by means of a unitary transformation with respect to the relative coordinates of all pairs

$$\mathbf{C} = \exp[-i\mathbf{G}] = \exp\left[-i\sum_{i < j} \mathbf{g}_{ij}\right]$$

$$\mathbf{G}^\dagger = \mathbf{G}$$
$$\mathbf{C}^\dagger \mathbf{C} = 1$$

## Correlated States

$$|\tilde{\psi}\rangle = \mathbf{C} |\psi\rangle$$

## Correlated Operators

$$\tilde{\mathbf{O}} = \mathbf{C}^\dagger \mathbf{O} \mathbf{C}$$

$$\langle \tilde{\psi} | \mathbf{O} | \tilde{\psi}' \rangle = \langle \psi | \mathbf{C}^\dagger \mathbf{O} \mathbf{C} | \psi' \rangle = \langle \psi | \tilde{\mathbf{O}} | \psi' \rangle$$

# Central and Tensor Correlators

$$\mathbf{C} = \mathbf{C}_\Omega \mathbf{C}_r$$

## Central Correlator $C_r$

- radial distance-dependent shift in the relative coordinate of a nucleon pair

$$g_r = \frac{1}{2} [s(r) \mathbf{q}_r + \mathbf{q}_r s(r)]$$

$$\mathbf{q}_r = \frac{1}{2} [\frac{\vec{r}}{r} \cdot \vec{q} + \vec{q} \cdot \frac{\vec{r}}{r}]$$

## Tensor Correlator $C_\Omega$



- angular shift depending on the orientation of spin and relative coordinate of a nucleon pair

$$g_\Omega = \frac{3}{2} \vartheta(r) [(\vec{\sigma}_1 \cdot \vec{q}_\Omega)(\vec{\sigma}_2 \cdot \vec{r}) + (\vec{r} \leftrightarrow \vec{q}_\Omega)]$$

$$\vec{q}_\Omega = \vec{q} - \frac{\vec{r}}{r} \mathbf{q}_r$$

$s(r)$  and  $\vartheta(r)$   
for given potential determined  
in the two-body system

# Correlated States: The Deuteron




# Correlated Interaction: $V_{\text{UCOM}}$

$$\tilde{\mathbf{H}} = \mathbf{T} + \mathbf{V}_{\text{UCOM}} + \mathbf{V}_{\text{UCOM}}^{[3]} + \dots$$

- **closed operator expression** for the correlated interaction  $\mathbf{V}_{\text{UCOM}}$  in two-body approximation
- correlated interaction and original NN-potential are **phase shift equivalent** by construction
- momentum-space matrix elements of correlated interaction are **similar to**  $V_{\text{low-}k}$
- operators of **all observables** (densities, transitions) have to be and can be **transformed consistently**

# Simplistic “Shell-Model” Calculation

- expectation value of Hamiltonian (with AV18) for Slater determinant of harmonic oscillator states



Application I



# No-Core Shell Model

in collaboration with  
Petr Navrátil (LLNL)



**No-Core Shell Model**  
+  
**Matrix Elements of Correlated  
Realistic NN-Interaction  $V_{\text{UCOM}}$**

- many-body state is expanded in Slater determinants of harmonic oscillator single-particle states
- large scale diagonalisation of Hamiltonian within a truncated model space ( $\mathcal{N}\hbar\omega$  truncation)
- assessment of short- and long-range correlations


# $^4\text{He}$ : Convergence



# $^4\text{He}$ : Convergence




# Tjon-Line and Correlator Range



- **Tjon-line**:  $E(^4\text{He})$  vs.  $E(^3\text{H})$  for phase-shift equivalent NN-interactions


# Tjon-Line and Correlator Range



- **Tjon-line:**  $E(^4\text{He})$  vs.  $E(^3\text{H})$  for phase-shift equivalent NN-interactions
- change of  $C_\Omega$ -correlator range results in shift along Tjon-line

**minimise net three-body force**  
by choosing correlator with energies close to experimental value



# Tjon-Line and Correlator Range



- **Tjon-line:**  $E(^4\text{He})$  vs.  $E(^3\text{H})$  for phase-shift equivalent NN-interactions
- change of  $C_\Omega$ -correlator range results in shift along Tjon-line


**minimise net  
three-body force**  
by choosing correlator  
with energies close to  
experimental value

# ${}^6\text{Li}$ : NCSM throughout the p-Shell




systematic NCSM studies  
throughout p-shell with  $V_{\text{UCOM}}$   
(+ Lee-Suzuki transformation)

# $^{10}\text{B}$ : Hallmark of a 3N Interaction?



# $^{10}\text{B}$ : Hallmark of a 3N Interaction?




# Outlook: NCSM beyond the p-Shell

## NCSM

- converged calculations essentially restricted to p-shell
- $6\hbar\omega$  calculation for  $^{40}\text{Ca}$  presently not feasible ( $\sim 10^{10}$  states)

## Importance Sampling NCSM

- diagonalization in space of **important** many-body configurations
- **a priori importance measure** given by perturbation theory




Application II:

# Hartree-Fock & Beyond


Standard Hartree-Fock  
+  
Matrix Elements of Correlated  
Realistic NN-Interaction  $V_{\text{UCOM}}$

- many-body state is a **Slater determinant** of single-particle states expanded in oscillator basis ( $\sim 13$  major shells)
- **correlations cannot be described** by Hartree-Fock states
- starting point for **improved many-body calculations**: MBPT, RPA, SM/CI, CC,...

# Hartree-Fock with VUCOM



# Perturbation Theory with V<sub>UCOM</sub>




# Application III

# RPA & Beyond


RPA, ERPA & SRPA  
+  
Matrix Elements of Correlated  
Realistic NN-Interaction  $V_{\text{UCOM}}$

- **fully self-consistent RPA** based on the Hartree-Fock orbits using the same  $V_{\text{UCOM}}$
- recovering sum rules with high precision, spurious center-of-mass mode fully decoupled at  $\sim 10$  keV
- **Extended-RPA and Second-RPA** to include effects of ground state correlations and complex configurations


# Isoscalar Giant Monopole



**prediction for ISGMR  
in good agreement with  
experiment**




**incompressibility  
resulting from  $V_{\text{UCOM}}$  is  
reasonable**




fm $^4$

# Isovector Dipole & Isoscalar Quadrupole



# Outlook: RPA with Three-Body Forces



- long-range tensor correlator & **repulsive three-body contact interaction**
  - systematic improvement of
    - rms-radii
    - single-particle spectra
    - strength distributions
- standard  $V_{\text{UCOM}}$   
—  $V_{\text{UCOM}}$  with long-range tensor & three-body contact interaction

Application IV

# Fermionic Molecular Dynamics (FMD)

# UCOM-FMD Approach

## Gaussian Single-Particle States

$$|\mathbf{q}\rangle = \sum_{\nu=1}^n \mathbf{c}_{\nu} |\mathbf{a}_{\nu}, \vec{\mathbf{b}}_{\nu}\rangle \otimes |\chi_{\nu}\rangle \otimes |m_t\rangle$$

$$\langle \vec{x} | \mathbf{a}_{\nu}, \vec{\mathbf{b}}_{\nu} \rangle = \exp \left[ - \frac{(\vec{x} - \vec{\mathbf{b}}_{\nu})^2}{2 \mathbf{a}_{\nu}} \right]$$

$\mathbf{a}_{\nu}$  : complex width

$\chi_{\nu}$  : spin orientation

$\vec{\mathbf{b}}_{\nu}$  : mean position & momentum

## Slater Determinant

$$|\mathbf{Q}\rangle = \mathcal{A} (|q_1\rangle \otimes |q_2\rangle \otimes \cdots \otimes |q_A\rangle)$$

## Correlated Hamiltonian

$$\tilde{\mathbf{H}} = \mathbf{T} + \mathbf{V}_{\text{UCOM}} + \delta V_{c+p+ls}$$

## Variation

$$\frac{\langle \mathbf{Q} | \tilde{\mathbf{H}} - \mathbf{T}_{\text{cm}} | \mathbf{Q} \rangle}{\langle \mathbf{Q} | \mathbf{Q} \rangle} \rightarrow \min$$


## Projection

restoration of rotational  
and inversion symmetry  
PAV / VAP


## Multi- Configuration

mixing of several  
intrinsic configurations  
GCM

# Intrinsic One-Body Density Distributions





# Structure of $^{12}\text{C}$




	$E$ [MeV]	$R_{ch}$ [fm]	$B(E2)$ [ $e^2 \text{ fm}^4$ ]
V/PAV	81.4	2.36	-
VAP $\alpha$ -cluster	79.1	2.70	76.9
PAV $^\pi$	88.5	2.51	36.3
VAP	89.2	2.42	26.8
Multi-Config	92.2	2.52	42.8
Experiment	92.2	2.47	$39.7 \pm 3.3$

# Structure of $^{12}\text{C}$ — Hoyle State




	Multi-Config	Experiment
$E$ [MeV]	92.4	92.2
$R_{\text{ch}}$ [fm]	2.52	2.47
$B(E2, 0_1^+ \rightarrow 2_1^+) [e^2 \text{ fm}^4]$	42.9	$39.7 \pm 3.3$
$M(E0, 0_1^+ \rightarrow 0_2^+) [\text{fm}^2]$	5.67	$5.5 \pm 0.2$

# Outlook: Resonances & Scattering in FMD



- collective coordinate representation as tool for the description of continuum states in FMD



first steps towards  
fully microscopic and  
consistent description  
of **structure and  
reactions**

# Conclusions

## ■ **Unitary Correlation Operator Method (UCOM)**

- explicit description of short-range central and tensor correlations
- universal phase-shift equivalent correlated interaction  $V_{UCOM}$

## ■ **Innovative Many-Body Methods**

- No-Core Shell Model
- Hartree-Fock, MBPT, SM/CI, CC, RPA, ERPA, SRPA,...
- Fermionic Molecular Dynamics

**unified description of nuclear  
structure across the whole  
nuclear chart is within reach**

# Epilogue

## ■ thanks to my group & my collaborators

- T. Böhlen, P. Hefeld, H. Hergert, M. Hild, N. Paar,  
P. Papakonstantinou, F. Schmitt, I. Türschmann, A. Zapp

Institut für Kernphysik, TU Darmstadt

- P. Navrátil

Lawrence Livermore National Laboratory

- T. Neff

NSCL, Michigan State University

- H. Feldmeier, C. Barbieri, S. Bacca, C. Özen,...

Gesellschaft für Schwerionenforschung (GSI)



supported by the DFG through SFB 634  
“Nuclear Structure, Nuclear Astrophysics and  
Fundamental Experiments...”