

The dipole strength distribution of ^8He and decay characteristics

C. Lehr,¹ T. Aumann,^{1,2,3} M. Duer,^{1,*} A. T. Saito,⁴ T. Nakamura,⁴ N. L. Achouri,⁵ D. Ahn,⁶ H. Baba,⁶ S. Bacca,⁷ C. A. Bertulani,⁸ M. Böhmer,⁹ F. Bonaiti,^{7,10,11} K. Boretzky,² C. Caesar,^{1,2,6} N. Chiga,⁶ D. Cortina-Gil,¹² C. A. Douma,¹³ F. Dufter,⁹ Z. Elekes,^{14,15} J. Feng,¹⁶ B. Fernández-Domínguez,¹² U. Forsberg,¹⁷ N. Fukuda,⁶ I. Gasparic,^{18,1,6} Z. Ge,⁶ R. Gernhäuser,⁹ J. M. Gheller,¹⁹ J. Gibelin,⁵ A. Gillibert,¹⁹ K. I. Hahn,^{20,21} Z. Halász,¹⁴ M. N. Harakeh,¹³ A. Hirayama,⁴ M. Holl,¹ N. Inabe,⁶ T. Isobe,⁶ J. Kahlbow,¹ N. Kalantar-Nayestanaki,¹³ D. Kim,²¹ S. Kim,^{21,1} T. Kobayashi,^{22,†} Y. Kondo,⁴ D. Körper,² P. Koseoglou,¹ Y. Kubota,⁶ P. J. Li,^{23,24} S. Lindberg,²⁵ Y. Liu,¹⁶ F. M. Marqués,⁵ S. Masuoka,²⁶ M. Matsumoto,⁴ J. Mayer,²⁷ K. Miki,^{1,22} M. Miwa,⁶ B. Monteagudo,⁵ A. Obertelli,^{1,19} N. A. Orr,⁵ H. Otsu,⁶ V. Panin,^{6,2} S. Y. Park,^{20,21} M. Parlog,⁵ S. Paschalis,^{17,1} P. M. Potlog,²⁸ S. Reichert,⁹ A. Revel,^{5,29,10,30} D. M. Rossi,¹ R. Roth,¹ M. Sasano,⁶ H. Scheit,¹ F. Schindler,¹ T. Shimada,⁴ S. Shimoura,²⁶ H. Simon,² S. Störck Dutine,¹ L. Stuhl,^{21,26} H. Suzuki,⁶ D. Symochko,¹ H. Takeda,⁶ S. Takeuchi,⁴ J. Tanaka,^{1,6} Y. Togano,⁴ T. Tomai,⁴ H. T. Törnqvist,^{1,2} J. Tscheuschner,¹ T. Uesaka,⁶ V. Wagner,¹ H. Yamada,⁴ B. Yang,¹⁶ L. Yang,²⁶ Z. H. Yang,⁶ M. Yasuda,⁴ K. Yoneda,⁶ L. Zanetti,¹ and J. Zenihiro,^{6,31}

¹ Technische Universität Darmstadt, Fachbereich Physik, 64289 Darmstadt, Germany

² GSI Helmholtzzentrum für Schwerionenforschung GmbH, Planckstr. 1, 64291 Darmstadt, Germany

³ Helmholtz Forschungsakademie Hessen für FAIR, Max-von-Laue-Str. 12, 60438 Frankfurt, Germany

⁴ Department of Physics, Institute of Science Tokyo, 2-12-1 O-Okayama, Meguro, Tokyo 152-8551, Japan

⁵ Université de Caen Normandie, ENSICAEN, IN2P3/CNRS, LPC-Caen UMR6534, F-14000 Caen, France

⁶ RIKEN Nishina Center for Accelerator-Based Science, 2-1 Hirosawa, Wako 351-0198, Japan

⁷ Institut für Kernphysik and PRISMA+ Cluster of Excellence, Johannes Gutenberg-Universität, 55128 Mainz, Germany

⁸ Department of Physics and Astronomy, East Texas A&M University, Commerce, TX 75429, USA

⁹ Technische Universität München, Physik Department, 85748 Garching, Germany

¹⁰ Facility for Rare Isotope Beams, Michigan State University, East Lansing, MI 48824, USA

¹¹ Physics Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA

¹² Dpt. de Física de Partículas, Universidade de Santiago de Compostela, E-15782 Santiago de Compostela, Spain

¹³ Nuclear Energy group, ESRIG, University of Groningen, 9747 AA Groningen, The Netherlands

¹⁴ HUN-REN ATOMKI, P.O. Box 51, H-4001 Debrecen, Hungary

¹⁵ Institute of Physics, Faculty of Science and Technology, University of Debrecen, Egyetem tér 1, H-4032 Debrecen, Hungary

¹⁶ State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

¹⁷ School of Physics, Engineering and Technology, University of York, York YO10 5DD

¹⁸ Rudjer Bošković Institute, Zagreb, Croatia

¹⁹ IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

²⁰ Ewha Womans University, Seoul 03760, Korea

²¹ Center for Exotic Nuclear Studies, Institute for Basic Science, Daejeon 34126, Korea

²² Department of Physics, Tohoku University, Miyagi 980-8578, Japan

²³ Department of Physics, The University of Hong Kong, Hong Kong, China

²⁴ Key Laboratory of Nuclear Physics and Ion-Beam Application (MOE),

Institute of Modern Physics, Fudan University, Shanghai 200433, China

²⁵ Department of Physics, Chalmers University of Technology, 412 96, Göteborg, Sweden

²⁶ Center for Nuclear Study, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

²⁷ Universität zu Köln, Institut für Kernphysik, Zùlpicher Straße 77, 50937 Köln, Germany

²⁸ Institute of Space Sciences, 077125 Magurele, Romania

²⁹ GANIL, CEA/DRF-CNRS/IN2P3, 14076 Caen, France

³⁰ Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA

³¹ Department of Physics, Kyoto University, Kitashirakawa-Oiwake, Sakyo, Kyoto 606-8502, Japan

(Dated: August 17, 2026)

The weak binding and spatially extended neutron densities characteristic of drip-line nuclei give rise to a distinctive low-energy dipole response. The drip-line nucleus ^8He is the most neutron-rich bound nucleus with a mass-to-charge ratio of $A/Z = 4$. We measure the dipole response of ^8He , including for the first time the four-neutron decay channel. A total dipole strength of $\sum B(E1)(E^* < 15 \text{ MeV}) = 0.95(16) \text{ e}^2\text{fm}^2$ and a dipole polarizability of $\alpha_D = 0.61(1) \text{ fm}^3$ are extracted from the differential Coulomb-excitation cross section and compared to state-of-the-art theoretical calculations employing coupled cluster and three-body approaches. We find that the dipole continuum is dominated, even at high excitation energies well above the $4n$ decay threshold, by two-neutron emission, pointing to a $^6\text{He}+2n$ structure of the excited dipole mode. No indication was found for a $4n$ final-state correlation, while pronounced nn and $^6\text{He}-n$ final-state correlations are apparent.

Introduction.— The dipole strength distributions of nuclei are a cornerstone of nuclear physics [1–3]. They

are essential for applications in nuclear astrophysics [4], medical isotope production [5], and fission [6]. At the same time, they provide key insights into nuclear structure and dynamics, especially at the limits of stability. At the neutron drip line, the weak binding of excess neutrons can lead to intriguing phenomena such as the development of neutron halos and thick neutron skins. These structural changes manifest in a characteristic manner in the electric dipole ($E1$) response. For stable nuclei, the $E1$ strength is dominated by the isovector giant dipole resonance (IVGDR), commonly understood as a collective harmonic vibration of neutrons versus protons. A unique feature of exotic nuclei is the appearance of significant $E1$ strength at lower excitation energies, known as soft $E1$ excitation, which is a direct consequence of the extended neutron densities. The mechanism of an enhanced $E1$ strength can be of non-resonant character, *i.e.*, a direct breakup leading to continuum states [7], as established for $1n$ halos, *e.g.*, ^{11}Be [8], or a resonance occurring as an oscillation of the core against the halo neutron(s) [9]. In $2n$ halos, *e.g.*, ^6He [10, 11] and ^{11}Li [12], the response is currently also understood as attributed to a non-resonant origin, while it is sensitive to ground-state correlations among the neutrons, as well as final-state interactions. Of particular interest is ^8He , the most neutron-rich bound nucleus, with an extreme mass-to-charge ratio of $A/Z = 4$, where the four excess neutrons are considered as forming a neutron-skin structure [13].

From the theoretical side, while the description of collective excitations has been traditionally dominated by density functional theory (DFT), especially in heavy-mass nuclei, first *ab initio* calculations recently became available for light- and medium-mass nuclei. A study of the ^8He dipole properties was carried out using coupled-cluster (CC) theory combined with the Lorentz integral transform (LIT) [14] in a LIT-CC approach, revealing a low-energy dipole strength around $E^* = 5$ MeV. Another study employing the importance-truncated no-core shell model (IT-NCSM) [15] likewise found a significant strength at low excitation energies. This is in contrast, *e.g.*, with a recent calculation based on the random-phase approximation (RPA) within the DFT framework [16], which disfavored the emergence of a soft dipole mode in ^8He , attributing the low-energy feature in the calculation to a spurious center of mass contamination. Experimental data can shed light on this controversy.

An established tool to experimentally study the $E1$ response of exotic nuclei is relativistic Coulomb excitation. In such a reaction, the projectile is excited by the Lorentz-contracted electromagnetic field of a high- Z target. If it is excited beyond its particle decay threshold, it will subsequently breakup. In ^8He , the lowest particle emission threshold is $^6\text{He}+2n$ with a two-neutron separation energy of $S_{2n} = 2.1$ MeV, while the four-neutron channel opens at an excitation energy of $S_{4n} = 3.1$ MeV. Coulomb excitation data for ^8He are

rather scarce, and in particular, only the two-neutron decay channel has been measured so far. A measurement performed at GSI at an incident beam energy of 227 MeV/nucleon [17], indicated the existence of a soft dipole resonance at $E^* \sim 4$ MeV. It was further supported via a transfer reaction measurement [18], although at a lower excitation energy of 3 MeV. On the other hand, an experiment carried out at MSU with a beam energy of 24 MeV/nucleon [19], attributed an insignificant fraction ($<3\%$) of the total energy-weighted dipole sum rule to a possible soft dipole mode. A recent inelastic proton-scattering experiment [20] confirmed this result, where the measured angular distribution was shown to be inconsistent with a dipole excitation. As both Coulomb dissociation experiments feature low statistics and measured only the $^6\text{He}+2n$ decay channel, a meaningful interpretation of the data and comparison to *ab initio* theory was difficult.

In this work, the high beam intensity provided at the Radioactive Ion Beam Factory (RIBF) in combination with a dedicated neutron detection setup, allowed to obtain a high-statistics data set with an extended excitation-energy range (up to 15 MeV) and to measure for the first time the $^4\text{He}+4n$ decay channel. Detection of four neutrons in coincidence is a challenging task, and only very recently the first successful measurement was demonstrated [21], within the same experimental campaign.

Experiment.— The experiment was performed at the RIBF operated by the RIKEN Nishina Center and the Center for Nuclear Study, University of Tokyo. The secondary beam of ^8He was produced at an energy of ~ 185 MeV/nucleon by fragmentation of a primary ^{18}O beam on a beryllium target, and selected using the BigRIPS fragment separator [22]. Projectiles were identified on an event-by-event basis via measurement of their energy deposition and time-of-flight (TOF) in thin plastic scintillators and transported to the SAMURAI spectrometer setup [23] with a typical ^8He intensity of $1.1 \cdot 10^5$ particles per second with 98% purity. Using two drift chambers, the projectiles were tracked towards the solid targets, where nuclear and Coulomb-excitation processes were induced. The main data were taken with a $3.157(47)$ g/cm² lead target. In addition, a set of targets with increasing Z was used, namely, carbon ($2.371(25)$ g/cm²), titanium ($3.115(49)$ g/cm²), and tin ($2.922(47)$ g/cm²). Data from these targets were used to correct for background from nuclear excitation reactions with high accuracy. An empty-target setting was used for additional background subtraction. The multi-neutron decay of ^8He following electromagnetic excitation was measured in complete kinematics via the $^6\text{He}+2n$ and $^4\text{He}+4n$ channels. The SAMURAI spectrometer, operated with a central magnetic field of 2.6 T, provided momentum analysis of the outgoing charged particles. Their trajectories were determined using two drift cham-

bers, placed before and after the dipole magnet, while scintillator walls at the focal plane were used for energy deposition and TOF measurements. Decay neutrons were measured at forward angles by three large-area segmented scintillator walls, consisting of the NeuLAND demonstrator [24] and the NEBULA array [25, 26]. The NeuLAND demonstrator, with an overall depth of 40 cm, was placed ~ 10.5 m downstream the target and separated by ~ 2 m from NEBULA. The latter was arranged in a two-wall configuration separated by 85 cm, with a depth of 2×24 cm. An additional thin plane of scintillators was placed in front of each wall to veto background from charged particles. The neutron momenta were derived from the TOF and the hit position measured in the detector arrays. A dedicated multi-neutron reconstruction algorithm based on Geant4 [27] simulations was developed, with special care to cross-talk rejection, in particular for the $4n$ case [28]. This resulted in total detection efficiencies of $\sim 1.2\%$ for the ${}^4\text{He}+4n$ channel, and $\sim 12\%$ for ${}^6\text{He}+2n$ at 1 MeV relative energy of the 5-body and 3-body systems, respectively. The simulations were validated as part of the experimental campaign by comparing the single-neutron detection efficiency [26]. To account for small deviations, a scaling factor is applied with an associated systematic uncertainty ($\sim 4\%$). For the cross-talk contribution, a 10% relative uncertainty is estimated. The decay energy of ${}^8\text{He}$ is reconstructed from the relative energy of the decay products (E_{fxn}) based on the invariant-mass method, using the four-momenta of the fragment (f) and x neutrons. Simulated resolutions of 0.2 and 0.24 MeV (σ) were obtained at 1 MeV relative energy for the two channels, respectively.

To demonstrate the neutron detection performance, the two-body decay of the ${}^7\text{He}$ ground-state resonance into ${}^6\text{He}+1n$ has been measured as a benchmark, populated via the one-neutron removal reaction ${}^8\text{He}(-1n)$ with a carbon target. The corresponding relative-energy (E_{fn}) spectrum was extracted and corrected for the detection efficiency to deduce the cross section. Following that, background contributions from empty-target data as well as remaining spurious events from the $2n$ breakup channels, which can be misidentified as $1n$ events, were subtracted. The ${}^7\text{He}$ spectrum is shown in Fig. 1, exhibiting the known $3/2^-$ ground-state resonance at $E_{fn} \sim 0.4$ MeV. The spectrum is fitted with an energy-dependent Breit-Wigner line shape convoluted with the experimental response to account for the relative-energy resolution (~ 90 keV (σ) at $E_{fn}=0.4$ MeV). The resonance energy and width, determined from the fit are $E_r = 385 \pm 1(\text{stat.}) \pm 1(\text{sys.})$ keV and $\Gamma = 193 \pm 1(\text{stat.}) \pm 4(\text{sys.})$ keV, respectively. These are in very good agreement with previously measured values using liquid-hydrogen target [29–31], but have been determined here with higher precision. The integrated cross section ($E_{fn} < 5$ MeV) for populating the ${}^7\text{He}$ resonant state is extracted as

$$185 \pm 1(\text{stat.}) \pm 15(\text{sys.}) \text{ mb.}$$




FIG. 1. Two-body relative-energy spectrum of the ${}^7\text{He}$ resonance populated by ${}^8\text{He}(-1n)$ removal reaction induced by a carbon target. The dark purple curve represents a fit to an energy-dependent Breit-Wigner line shape, convoluted with the experimental response. The dotted-dashed curve displays the overall detection efficiency.

Results.— The ${}^6\text{He}+2n$ and ${}^4\text{He}+4n$ breakup channels of ${}^8\text{He}$ induced by a lead target were identified by selecting the outgoing fragment in coincidence with 2 and 4 neutrons, respectively. The three- and five-body relative-energy spectra are then reconstructed via the invariant mass following subtraction of background contributions as determined from the empty-target measurement. For cross-talk correction, the corresponding contributions for the two channels are considered and subtracted (see Supplemental Material [32]). Finally, the measured spectra are corrected for detection efficiency to determine the differential cross section.

To extract pure electromagnetic properties and the dipole response of ${}^8\text{He}$, the nuclear part of the cross section has to be eliminated. This is often done by a measurement with a light target such as carbon [33], where the electromagnetic excitation cross section is small. Under the assumption that the nuclear spectra have the same shape when impinging on both target nuclei, the obtained cross section is then scaled and subtracted from the total cross section measured with the lead target. The scaling factor is typically obtained semi-empirically [34], or by a theoretical calculation [10]. In this work, it was determined independently based on measurements with a series of targets, thus minimizing model dependence. While the Coulomb excitation cross section scales approximately as Z^2 , the nuclear cross section for peripheral reactions relevant for halo nuclei scales as $A^{1/3}$ [33]. This dependency is used to extract the scaling factor from the data by assuring that the extracted $B(E1)$ distribution is independent from the target (Sn, Pb) by introducing a free parameter for the scaling and applying a fitting

procedure. Finally, the small electromagnetic contribution within the carbon-target spectrum is corrected for (see Supplemental Material [32]).




FIG. 2. Coulomb excitation cross section spectra for three- (dark green) and five-body (light green) breakup channels of ^8He measured with lead target. The inset shows the corresponding dipole strength distributions, $dB(E1)/dE^*$, as a function of the excitation energy.

The differential Coulomb excitation cross sections for the $2n$ and $4n$ channels are shown in Fig. 2. The integrated cross sections up to $E^* = 15$ MeV are $\sigma_{C,2n} = 243 \pm 1(\text{stat.}) \pm 33(\text{sys.})$ mb and $\sigma_{C,4n} = 21.14 \pm 0.02(\text{stat.}) \pm 5.34(\text{sys.})$ mb. Systematic errors include uncertainties of the detection efficiency, target areal densities and the scaling factor for the subtraction of the nuclear contribution. From the differential cross section, the dipole strength distributions $dB(E1)/dE^*$ are derived following Ref. [35]. The excitation process is treated as a photo-absorption induced by a virtual photon, known as the equivalent-photon method. The Coulomb breakup cross section for electric dipole excitation is expressed as

$$\frac{d\sigma(E1)}{dE^*} = \frac{16\pi^3}{9\hbar c} N_{E1}(E^*) \frac{dB(E1)}{dE^*}, \quad (1)$$

where $N_{E1}(E^*)$ is the number of virtual photons at excitation energy E^* . This relation connects the measured cross section to the reduced transition probability $B(E1)$, which contains the physics information of the dipole response (see Fig. 2 inset). It can be clearly observed that the $2n$ channel is dominant even at high excitation energies well above the $4n$ threshold. This surprising result points towards a characteristic structure effect of the ^8He dipole continuum. The dipole vibration seems to be dominated by a ^6He -di-neutron component, which would explain the preferred decay into $^6\text{He}+2n$.

The total $dB(E1)/dE^*$ spectrum, (sum of the $2n$ and $4n$ channels) is shown in Fig. 3. From that,




FIG. 3. Total $B(E1)$ spectrum measured with lead target. Theory predictions folded with the experimental response are shown in blue based on CC [14] (band), HH [37] (solid curve) and COSM [38] (dotted-dashed curve).

the non-energy-weighted dipole strength integrated up to $E^*=15$ MeV amounts to $\Sigma B(E1) = 0.95 \pm 0.01(\text{stat.}) \pm 0.16(\text{sys.}) e^2\text{fm}^2$ and the energy-weighted one to $\Sigma E^* B(E1) = 7.55 \pm 0.18(\text{stat.}) \pm 1.39(\text{sys.}) e^2\text{fm}^2\text{MeV}$, which corresponds to about 1/3 of the Thomas-Reiche-Kuhn sum rule [36]. With the inverse energy-weighted sum of the dipole strength, the electric dipole polarizability is extracted as

$$\alpha_D = \frac{8\pi}{9} \int \frac{dB(E1)/dE^*}{E^*} dE^*, \quad (2)$$

resulting in $\alpha_D = 0.61 \pm 0.01(\text{stat.}) \pm 0.01(\text{sys.}) \text{fm}^3$.

The $B(E1)$ distribution exhibits a clear soft-dipole enhancement at 3 MeV. This is consistent with previous experiments [17, 18] where only the $2n$ channel was measured. The weak contribution of the $4n$ channel to the total strength confirms it is not caused by this channel opening at around this energy, thus resolving discussions on the interpretation of the peak observed in earlier experiments.

We compare the data with ab initio predictions from the LIT-CC method [14, 39]. Refs. [14, 39] focused on moments of the dipole response function of ^8He , and in particular on α_D , which was found to vary between 0.37(3) to 0.48(3) fm^3 employing different chiral effective field theory interactions. In the present work, we determine the full dipole response function, $dB(E1)/dE^*$, by numerically inverting the Lorentz integral transform. We focus on the chiral interaction 1.8/2.0 (EM) [40], which provides the largest α_D . LIT-CC calculations are performed with the best approximation scheme at disposal, labeled CCSDT-1, including linear triples corrections in both the ground- and excited-state calculations [41]. The resulting response function is shown in

Fig. 3 with a blue band quantifying the uncertainties associated with the inversion procedure and model-space convergence. Further details on the LIT-CC calculations are provided in the Supplemental Material [32]. While it describes the strength distribution rather well at excitation energies above 5-6 MeV, it does not reproduce the low-energy part of the spectrum which peaks at ~ 3 MeV. A possible origin for this discrepancy may lie in missing higher-order correlations within the CC many-body expansion. To explore this, we compare LIT-CC calculations with results from the IT-NCSM approach [15], which is expected to capture a larger portion of many-body correlations for the same interaction. Similar to the LIT-CC case, the spread in α_D predictions for different chiral forces remains significant, ranging from 0.4454(19) to 0.5950(1) fm³. For the NNLO_{sat} interaction [42], where both results are available, IT-NCSM predicts $\alpha_D = 0.4454(19)$ fm³, which is more than 10% higher than the corresponding LIT-CC value of 0.37(3) fm³, suggesting that many-body correlations beyond triples might play a role for this nucleus. Since only a discrete strength distribution is calculated in this case, it cannot be compared directly to the data.

The strength function for the $2n$ breakup channel was also estimated in a three-body model and solving the Schrödinger equation with a source term (dipole operator acting on the ground-state wave function) using the hyperspherical harmonics (HH) method [37] (solid curve). The rising edge of the peak at $E^* \sim 3$ MeV is reproduced correctly, while for higher energies a large discrepancy is visible, despite the fact that the measured spectrum is still dominated by the $2n$ decay. Finally, a five-body ($^4\text{He}+4n$) cluster orbital shell model (COSM) was employed using the complex-scaling method to describe many-body unbound states of ^8He [38]. Different excitation modes were found to coexist in the dipole strength distribution (dotted-dashed curve) - below 10 MeV the strength comes from single-particle excitation, $^7\text{He}+n$, leading to sequential decay into $^6\text{He}+2n$. At around 13 MeV, collective excitations of four neutrons with strong configuration mixing are enhanced, indicating a possible soft dipole mode.

Correlations in the final state can be investigated by considering the two-body sub-systems. For the $^6\text{He}+2n$ decay channel this corresponds to the neutron-neutron (nn) and fragment-neutron (fn) and their corresponding two-body relative energies E_{nn} and E_{fn} . For the latter, two possible combinations are taken into account with either of the two neutrons. The two-body correlations are revealed by dividing the measured spectra by simulated ones. Simulations of the three-body ($^6\text{He}+2n$) system were performed with the differential cross section spectrum as an input, and modeling the breakup after the reaction using a phase-space decay. The measured and simulated data are normalized to each other, such that their division shows deviations of the experimental




FIG. 4. (a) Fragment-neutron (fn) two-body relative energy (E_{fn}) for $^8\text{He} \rightarrow ^6\text{He} + 2n$ breakup channel. Shown is the measured spectrum divided by a simulated one assuming phase-space decay. (b) Same for the neutron-neutron sub-system.

data from a pure phase-space decay.

Figure 4 (a) shows the E_{fn} relative-energy spectrum, with a clear enhancement at the ground-state energy of the ^7He resonance. This corresponds to the sequential decay of the three-body system through the resonant state, thus deviating from a direct phase-space decay. In the nn sub-system (panel (b)), low relative energies are enhanced as a result of the nn interaction, reflecting the di-neutron virtual state.

For the $^4\text{He}+4n$ decay channel, we study the $4n$ sub-system to get insight on correlations of the four neutrons. This is of particular interest, following the recent observation of a near-threshold (~ 2 MeV) structure in the $4n$ system via an α -knockout reaction from ^8He [43]. Figure 5 (top panel) shows the $4n$ relative-energy spectrum (E_{4n}) in comparison with a simulated phase-space decay, where the bottom panel presents the ratio between them. No indication for a $4n$ final-state interaction is seen. This is most likely related to the fact that the $4n$ decay sets in only at rather high excitation energy, as discussed above, preventing a pronounced $4n$ low-energy correlation. The phase-space decay seems to overall describe the data rather well, while deviations at low relative energy may hint of nn correlations.

Summary and conclusion.— In this Letter, we present the dipole response of the most neutron-rich bound nucleus ^8He up to an excitation energy of 15 MeV as extracted from a measurement of its Coulomb excitation cross section at an energy of ~ 185 MeV/nucleon. The excitation spectrum has been reconstructed from an exclusive measurement of the residues in the final state using the invariant-mass method, including for the first time besides the $2n$ channel the 5-body decay into $^4\text{He}+4n$. The dipole strength distribution shows a peak at around 3 MeV followed by a broad continuum up to 15 MeV. The dominant decay channel is thereby the $^6\text{He}+2n$ channel




FIG. 5. Top: $4n$ relative-energy spectrum (E_{4n}) for ${}^8\text{He} \rightarrow {}^4\text{He} + 4n$ breakup channel. The red curve represents a simulated distribution assuming 5-body phase-space decay. Bottom: measured E_{4n} distribution divided by the simulated one.

surprisingly even for excitation energies well above the $4n$ threshold, pointing to a dominant di-neutron correlation in the dipole continuum structure. This is supported by the observation of the 2-body correlations in the decay, where strong correlations at the ${}^7\text{He}$ resonance energy and the nn virtual state at low energy are observed. In contrast, no strong correlation among the 4 neutrons in the $4n$ channel has been observed, likely related to the fact that the $4n$ channel covers relatively large excitation energies.

The experimental dipole strength distribution has been compared to an ab initio LIT-CC calculation and to a phenomenological three-body model. The ab initio theory shows good agreement with experiment for excitation energies above 5 MeV, but fails to reproduce the soft-dipole peak at 3 MeV, while the phenomenological three-body model reproduces well the low-energy peak, but misses the higher-lying dipole strength. For the former, a possible reason may lie in missing correlations, while one does not expect the phenomenological cluster model to be valid in the region where the ${}^4\text{He}$ core is excited, which is well explained by the ab initio theory. More theoretical efforts need to be devoted in the future to fully describe the dipole strength distribution of extremely neutron-rich drip-line nuclei.

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)

through Project-ID 279384907 - SFB 1245 and Project ID 39083149 PRISMA⁺ EXC 2118/1, the GSI-TU Darmstadt cooperation agreement, by the German Federal Ministry of Education and Research - BMBF project number 05P15RDFN1. This work was also supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under the FRIB Theory Alliance award DE-SC0013617; by the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research and Office of Nuclear Physics, Scientific Discovery through Advanced Computing (SciDAC) program (SciDAC-5 NUCLEI). C.A. Bertulani acknowledges support from the U.S. Department of Energy, Office of Science, award DE-SC0026074. N.L.A., F.D., J. G., F. M. M., M.P. and N. A. O. acknowledge partial support from the Franco-Japanese LIA-International Associated Laboratory for Nuclear Structure Problems as well as the French ANR-14-CE33-0022-02 EXPAND. S.C.P. acknowledges support by UK STFC under contract numbers ST/P003885/1 and ST/L005727/1 and the University of York Pump Priming Fund. This work was partially supported by the Institute for Basic Science (IBS-R031-D1) in Korea. P. L. acknowledges support from the National Natural Science Foundation of China (Grant No. 12405147). T. N. and Y. K. acknowledge support from JSPS KAKENHI Grants No. JP24H00006 and No. JP18H05404. F.B. and S.B. thank G. Hagen for providing access to the nuclear coupled-cluster code developed at ORNL (NUCCOR).

This manuscript has been authored in part by UT-Battelle, LLC, under contract DE-AC05-00OR22725 with the US Department of Energy (DOE). The US government retains and the publisher, by accepting the article for publication, acknowledges that the US government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for US government purposes. DOE will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan (<http://energy.gov/downloads/doe-public-access-plan>).

* Corresponding Author: Meytal Duer (mduer@ikp.tu-darmstadt.de)

† Deceased

- [1] T. Aumann and T. Nakamura. The electric dipole response of exotic nuclei. *Phys. Scr.*, T152:014012, 2013.
- [2] S. Bacca and P. Saori. Electromagnetic reactions on light nuclei. *J. Phys. G: Nucl. Part. Phys.*, 41(12):123002, 2014.
- [3] M. N. Harakeh and A. van der Woude. *Giant Resonances*. Oxford Studies in Nuclear Physics. Oxford University Press, Oxford, 2001.
- [4] M. Arnould, S. Goriely, and K. Takahashi. The r-process of stellar nucleosynthesis: Astrophysics and nu-

- clear physics achievements and mysteries. *Phys. Rep.*, 450(4–6):97–213, September 2007.
- [5] Y. Wang, D. Chen, R. dos Santos Augusto, et al. Production review of accelerator-based medical isotopes. *Molecules*, 27(16):5294, 2022.
- [6] M.B. Chadwick et al. Endf/b-vii.1 nuclear data for science and technology: Cross sections, covariances, fission product yields and decay data. *Nucl. Data Sheets*, 112(12):2887–2996, 2011.
- [7] P.G. Hansen and B. Jonson. The Neutron Halo of Extremely Neutron-Rich Nuclei. *Europhys. Lett.*, 4:409, 1987.
- [8] T. Nakamura et al. Coulomb dissociation of a halo nucleus ^{11}Be at 72 AMeV. *Phys. Lett. B*, 331:296–301, 1994.
- [9] K. Ikeda. Structure of neutron rich nuclei. *Nucl. Phys. A*, 538:355c, 1992.
- [10] T. Aumann et al. Continuum excitations in ^6He . *Phys. Rev. C*, 59:1252–1262, 1999.
- [11] Y.L. Sun et al. Three-body breakup of ^6He and its halo structure. *Phys. Lett. B*, 814:136072, 2021.
- [12] T. Nakamura et al. Observation of Strong Low-Lying $E1$ Strength in the Two-Neutron Halo Nucleus ^{11}Li . *Phys. Rev. Lett.*, 96:252502, 2006.
- [13] I. Tanihata et al. Revelation of thick neutron skins in nuclei. *Phys. Lett. B*, 289:261, 1992.
- [14] F. Bonaiti, S. Bacca, and G. Hagen. Ab initio coupled-cluster calculations of ground and dipole excited states in ^8He . *Phys. Rev. C*, 105:034313, 2022.
- [15] C. Stumpf. *Nuclear Spectra and Strength Distributions from Importance-Truncated Configuration-Interaction Methods*. PhD thesis, Technische Universität Darmstadt, 2018.
- [16] J. Piekarewicz. Insights into the possible existence of a soft dipole mode in ^8He . *Phys. Rev. C*, 105:044310, 2022.
- [17] M. Meister, K. Markenroth, D. Aleksandrov, T. Aumann, T. Baumann, et al. ^8He – ^6He : a comparative study of electromagnetic fragmentation reactions. *Nucl. Phys. A*, 700:3, 2002.
- [18] M. S. Golovkov et al. The ^8He and ^{10}He spectra studied in the (t,p) reaction. *Phys. Lett. B*, 672:22–29, 2009.
- [19] Y. Iwata, K. Ieki, A. Galonsky, J. J. Kruse, J. Wang, et al. Dissociation of ^8He . *Phys. Rev. C*, 62:064311, 2000.
- [20] M. Holl et al. Proton inelastic scattering reveals deformation in ^8He . *Phys. Lett. B*, 822:136710, 2021.
- [21] Y. Kondo, N. L. Achouri, H. Al Falou, L. Atar, T. Aumann, et al. First observation of ^{28}O . *Nature*, 620:965, 2023.
- [22] T. Kubo et al. Status and Overview of Superconducting Radioactive Isotope Beam Separator BigRIPS at RIKEN. *IEEE Trans. Appl. Supercond.*, 17:1069–1077, 2007.
- [23] T. Kobayashi, N. Chiga, T. Isobe, Y. Kondo, T. Kubo, K. Kusaka, T. Motobayashi, T. Nakamura, J. Ohnishi, H. Okuno, et al. SAMURAI spectrometer for RI beam experiments. *Nucl. Instrum. Methods Phys. Res. Sect. B*, 317:294–304, 2013.
- [24] K. Boretzky et al. NeuLAND: The high-resolution neutron time-of-flight spectrometer for R^3B at FAIR. *Nucl. Instrum. Methods Phys. Res. Sect. A*, 1014:165701, 2021.
- [25] T. Nakamura and Y. Kondo. Large acceptance spectrometers for invariant mass spectroscopy of exotic nuclei and future developments. *Nucl. Inst. Meth. B*, 376:151, 2016.
- [26] Y. Kondo, T. Tomai, and T. Nakamura. Recent progress and developments for experimental studies with the SAMURAI spectrometer. *Nucl. Inst. Meth. B*, 463:173, 2020.
- [27] S. Agostinelli et al. Geant4—a simulation toolkit. *Nucl. Instrum. Methods Phys. Res. Sect. A*, 506:250, 2003.
- [28] C. Lehr. *Low-energy dipole response of the halo nuclei $^{6,8}\text{He}$* . PhD thesis, Technische Universität Darmstadt, 2021.
- [29] Yu. Aksyutina et al. Properties of the ^7He ground state from ^8He neutron knockout. *Phys. Lett. B*, 679:191–196, 2009.
- [30] D. H. Denby et al. Ground state energy and width of ^7He from ^8Li proton knockout. *Phys. Rev. C*, 78:044303, 2008.
- [31] F. Renzi et al. Spectroscopy of ^7He using the $^9\text{Be}(^6\text{He}, ^8\text{Be})$ transfer reaction. *Phys. Rev. C*, 94:024619, 2016.
- [32] See Supplemental Material <https://doi.org/10.1103/vp3m-5rpp> which includes Refs. [? ? ? ?] for LIT-CC calculations, cross-talk correction and nuclear background contribution.
- [33] T. Aumann. Low-energy dipole response of exotic nuclei. *Eur. Phys. J. A*, 55:234, 2019.
- [34] D. M. Rossi et al. Measurement of the Dipole Polarizability of the Unstable Neutron-Rich Nucleus ^{68}Ni . *Phys. Rev. Lett.*, 111:242503, 2013.
- [35] C. A. Bertulani and G. Baur. Electromagnetic processes in relativistic heavy ion collisions. *Phys. Rep.*, 163:299–408, 1988.
- [36] H. Sagawa and M. Honma. Sum rule approach to decoupled giant dipole state. *Phys. Lett. B*, 251:17, 1990.
- [37] L. V. Grigorenko et al. Soft Dipole Mode in $^8\text{He}^*$. *Phys. Part. Nucl. Lett.*, 6:118–125, 2009.
- [38] T. Myo and K. Katō. Modeling the electric dipole strength of neutron-rich ^8He : Prediction of multineutron collective excitations. *Phys. Rev. C*, 106:L021302, 2022.
- [39] F. Bonaiti and S. Bacca. Low-energy dipole strength in ^8He . *Few-Body Systems*, 65(2):54, 2024.
- [40] K. Hebeler, S. K. Bogner, R. J. Furnstahl, A. Nogga, and A. Schwenk. Improved nuclear matter calculations from chiral low-momentum interactions. *Phys. Rev. C*, 83:031301(R), 2011.
- [41] M. Miorrelli, S. Bacca, G. Hagen, and T. Papenbrock. Computing the dipole polarizability of ^{48}Ca with increased precision. *Phys. Rev. C*, 98:014324, Jul 2018.
- [42] A. Ekström et al. Accurate nuclear radii and binding energies from a chiral interaction. *Phys. Rev. C*, 91:051301, 2015.
- [43] M. Duer et al. Observation of a correlated free four-neutron system. *Nature*, 606:678, 2022.